



User Guide

Due to the pandemic's effect on the transplant system, adjustments have been made for this data reporting cycle and potentially future reporting cycles as well. For transplant programs, this means that SRTR will not include patient follow-up starting from March 12, 2020, the day prior to the emergency declaration, i.e., waitlist survival, transplant rate, and outcomes will not be assessed after that date.

This report contains a wide range of useful information about the kidney transplant program at New York University Medical Center (NYUC). The report has three main sections:

- A. Program Summary
- B. Waiting List Information
- C. Transplant Information

The Program Summary is a one-page summary highlighting characteristics of the program, including the number of candidates on the waiting list, the number of transplants performed at the program, the number of patients being cared for by the program, and patient outcomes, including outcomes while on the waiting list (the transplant rate and the death rate while on the waiting list) and outcomes after transplant (patient and graft survival probabilities). If the program performed transplants in both adults and children, survival probabilities for adults and children (pediatrics) are provided separately. For each of the outcomes measures presented, a comparison is provided showing what would be expected at this program if it were performing as similar programs around the country perform when treating similar patients. More details regarding these outcome measures are provided in Sections B and C of the report.

The Waiting List Information section contains more detailed information on how many candidates are on the waiting list at the program, the types of candidates on the waiting list, how long candidates typically have to wait for a transplant at this program, how frequently candidates successfully receive a transplant, and how often candidates on the waiting list die before receiving a transplant.

Table B1 shows the activity on this program's waiting list during two recent 1-year periods and provides comparisons to all programs within this program's OPTN region (see <http://optn.transplant.hrsa.gov/members/regions.asp> for information on OPTN regions) and the nation as a whole. Tables B2 and B3 describe the candidates on the waiting list at this program, with comparisons to candidates waiting in the same donor service area (OPO/DSA) the OPTN region, and the nation as a whole.

Table B4 shows how many candidates were removed from the waiting list because they received a transplant. The program's transplant rate is calculated as the number of candidates who received a transplant divided by the person-years observed at the program (person-years is a combination of how many candidates were on the waiting list along with how long each candidate was followed since some candidates are not on the waiting list for the entire year). The transplant rate and comparisons to what would be expected at this program are presented in Figures B1 and B2. Figure B1 shows the transplant rate compared to what was expected at this program. The expected transplant rate is an estimate of what we would expect at this program if it were performing transplants at rates similar to other programs in the US with similar candidates on their waiting lists. The expected rate is only an estimate, and is made with a certain level of uncertainty. This uncertainty is shown in Figure B2. Figure B2 displays the ratio of the observed to the expected transplant rate. A ratio of 1 indicates that the



User Guide

observed transplant rate was equal to the expected transplant rate, while a ratio less than 1 indicates the observed rate was lower than expected rate and a ratio greater than 1 indicates the observed rate was higher than the expected rate. However, the level of uncertainty must be considered when interpreting these numbers. The 95% interval is also shown on Figure B2. This interval provides a range within which the true ratio of observed to expected transplant rates is likely to be. If this confidence interval includes (crosses) 1.0, then we cannot say that this program's observed transplant rate is different from what would be expected. The observed transplant rate at this program was 47.1 per 100 person-years. Transplant rates are also provided for adult and pediatric patients separately along with comparisons to adult and pediatric rates in the DSA, the OPTN region, and the nation. Transplant rates are also presented excluding transplants from a living donor (Table B4D and Figures B1D-B3D). Please refer to the PSR Technical Methods documentation available at <http://www.srtr.org> for more detail regarding how expected rates are calculated.

The death rate (also known as the mortality rate) for candidates on the waiting list is presented in Table B5 and Figures B4-B6. These data are presented in the same way as the transplant rate data in the previous section. The intent of this table and figures is to describe risk of death once candidates are listed rather than while they are listed, but before they are transplanted. Therefore, time at risk and deaths after removal from the waiting list for reasons other than transplant, transfer to another transplant program, or recovery (no longer needing a transplant), and before any subsequent transplant, are included. As with transplant rates, mortality rates should be interpreted carefully taking into consideration the interval displayed in Figure B5. For a complete description of how observed and expected mortality rates are calculated, please refer to the technical documentation available at <http://www.srtr.org>.

Survival from listing is presented in Table B6 and Figures B7-B9. These data are presented in the same way as the waiting list mortality rate data in the previous section. The intent of this table and figures is to describe risk of death once candidates are listed rather than while they are listed, including after a transplant. As with transplant rates, mortality rates should be interpreted carefully taking into consideration the interval displayed in Figure B8. For a complete description of how observed and expected mortality rates are calculated, please refer to the technical documentation available at <http://www.srtr.org>.

Table B7 presents information on what happens to candidates on the waiting list by three different time points after listing: 6 months, 12 months, and 18 months. The table displays percentages of candidates who have died, been removed from the waiting list, been transplanted, or been transferred or lost-to-follow-up. Tables B8 and B9 provide more detail regarding how many candidates have received a deceased donor transplant by certain time points during the first 3 years after being put on the transplant waiting list. Each row of Tables B8 and B9 presents the percent of candidates who received a deceased donor transplant by each time point. Table B10 presents data on the time it took for different percentages of patients to be transplanted for candidates added to the list between 07/01/2014 and 12/31/2019. The time it took for 5% (the 5th percentile) of patients to receive a transplant at this program was 0.4 months. If "Not Observed" is displayed in the table, then too few candidates received transplants before 06/30/2020 to calculate a particular percentile of transplant times.



User Guide

Table B11 contains a summary of the offer acceptance practices of the program. The offer acceptance ratio indicates whether the program is more or less likely to accept offers than the average program. If the offer acceptance ratio is greater than 1.0, then the program tends to accept more offers than average; if the offer acceptance ratio is less than 1.0, then the program tends to accept fewer offers than average. Figure B10 shows the distribution of program offer acceptance rates as well as the offer acceptance rate for this program. Figures B11 - B14 similarly show offer acceptance rates for subsets of offers.

The Transplant Information section begins with descriptions of transplant recipients in Tables C1 and C2. Data on recipients of deceased donor transplants are presented (Tables C1D and C2D); if applicable, data on recipients of living donor transplants are presented separately (Tables C1L and C2L). Comparisons to the region and the nation as a whole are provided. A description of the deceased donors used at this program is provided in Table C3D, along with characteristics of living donors in Table C3L, if applicable. Finally, information on the transplant procedure for deceased and living donor transplants is presented in Tables C4D and C4L, respectively.

Starting with Table C5, transplant outcomes are presented along with comparisons to what would be expected at this program and what happened in the nation as a whole. Tables C5-C10 present information on graft survival (survival of the transplanted organ), with data presented separately for adult and pediatric recipients. Patients are followed from the time of transplant until either failure of the transplanted organ or death, whichever comes first. Please refer to the technical methods for more information on these calculations (<http://www.srtr.org>).

While Tables C5-C10 present data on graft survival, Tables C11-C16 present information on patient survival. For these tables, patients are followed from the time of transplant until death, regardless of whether the transplant is functioning or the patient required another transplant to survive.

Tables C17 and C18 summarize the multiorgan transplant outcomes at this program. The summary statistics in these tables are descriptive and are not risk-adjusted for different donor and candidate characteristics.

Table D1 shows the rates of follow-up for living donors.

Additional information regarding the technical methods and the risk adjustment models used to estimate expected event rates is available on the SRTR website at <http://www.srtr.org>. We welcome and encourage feedback on these reports. Please feel free to share feedback with the SRTR at the following e-mail: srtr@srtr.org.



Table of Contents

Section	Page
User Guide	i
A. Program Summary	
Program Summary	1
B. Waiting List Information	
Waiting list activity	2
Demographic characteristics of waiting list candidates	3
Medical characteristics of waiting list candidates	4
Transplant rates	5
Deceased donor transplant rates	6
Waiting list mortality rates	7
Patient survival from listing	8
Waiting list candidate status after listing	9
Percent of candidates with deceased donor transplants: demographic characteristics	10
Percent of candidates with deceased donor transplants: medical characteristics	11
Time to transplant for waiting list candidates	12
Offer acceptance practices	13
C. Transplant Information	
Deceased donor transplant recipient demographic characteristics	15
Living donor transplant recipient demographic characteristics	16
Deceased donor transplant recipient medical characteristics	17
Living donor transplant recipient medical characteristics	18
Deceased donor characteristics	19
Living donor characteristics	20
Deceased donor transplant characteristics	21
Living donor transplant characteristics	22
Graft survival	23
Patient survival	41
Multi-organ transplant graft survival	59
Multi-organ transplant patient survival	59
D. Living Donor Information	
Living donor follow-up summary	60



A. Program Summary

Figure A1. Waiting list and transplant activity




Table A1. Census of transplant recipients

Recipients	07/01/2018-06/30/2019	07/01/2019-06/30/2020
Transplanted at this center	196	186
Followed by this center*	456	587
...transplanted at this program	452	581
...transplanted elsewhere	4	6

* Recipients followed are transplant recipients for whom the center has submitted a post-transplant follow-up form for a transplant that took place before the 12-month interval for each column.

Figure A2. Transplant rates 07/01/2018 - 03/12/2020




Figure A3. Waiting list mortality rates 07/01/2018 - 03/12/2020




Figure A4. First-year adult graft and patient survival: 07/01/2017 - 12/31/2019





Figure A5. First-year pediatric graft and patient survival: 07/01/2017 - 12/31/2019





B. Waiting List Information

Table B1. Waiting list activity summary: 07/01/2018 - 06/30/2020

Waiting List Registrations	Counts for this center		Activity for 07/01/2019 to 06/30/2020 as percent of registrants on waiting list on 07/01/2019		
	07/01/2018-06/30/2019	07/01/2019-06/30/2020	This Center (%)	OPTN Region (%)	U.S. (%)
On waiting list at start	344	457	100.0	100.0	100.0
Additions					
New listings at this center	380	306	67.0	35.2	39.3
Removals					
Transferred to another center	6	6	1.3	2.4	1.7
Received living donor transplant*	56	50	10.9	6.3	5.7
Received deceased donor transplant*	139	136	29.8	12.2	16.6
Died	24	20	4.4	5.9	4.1
Transplanted at another center	7	15	3.3	2.5	3.8
Deteriorated	7	14	3.1	3.3	4.0
Recovered	0	0	0.0	0.1	0.2
Other reasons	28	18	3.9	4.3	5.0
On waiting list at end of period	457	504	110.3	98.2	98.2

* These patients were removed from waiting list with removal code indicating transplant; this may not equal the number of transplants performed at this center during the specified period.



B. Waiting List Information

Table B2. Demographic characteristics of waiting list candidates
Candidates registered on the waiting list between 07/01/2019 and 06/30/2020

Demographic Characteristic	New Waiting List Registrations 07/01/2019 to 06/30/2020 (%)			All Waiting List Registrations on 06/30/2020 (%)		
	This Center (N=306)	OPTN Region (N=2,751)	U.S. (N=39,776)	This Center (N=504)	OPTN Region (N=7,678)	U.S. (N=99,301)
All (%)	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity/Race (%)*						
White	33.0	39.6	41.7	28.4	30.1	35.2
African-American	33.0	30.7	28.5	31.5	36.2	32.1
Hispanic/Latino	16.3	17.8	19.7	21.0	20.6	21.0
Asian	17.3	10.7	8.1	19.0	12.2	9.9
Other	0.3	1.2	1.9	0.0	0.8	1.8
Unknown	0.0	0.0	0.0	0.0	0.0	0.0
Age (%)						
<2 years	0.0	0.1	0.1	0.0	0.1	0.1
2-11 years	0.0	0.6	0.9	0.0	0.5	0.6
12-17 years	1.3	1.3	1.5	0.8	1.4	1.0
18-34 years	13.4	11.4	10.7	13.7	10.0	10.3
35-49 years	24.5	21.6	24.2	26.8	24.7	26.9
50-64 years	38.9	42.1	41.3	41.1	44.1	43.4
65-69 years	10.8	12.5	13.3	8.7	11.5	12.1
70+ years	11.1	10.5	8.1	8.9	7.7	5.6
Gender (%)						
Male	65.7	63.8	62.1	61.5	62.5	62.1
Female	34.3	36.2	37.9	38.5	37.5	37.9

* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



B. Waiting List Information

Table B3. Medical characteristics of waiting list candidates
Candidates registered on the waiting list between 07/01/2019 and 06/30/2020

Medical Characteristic	New Waiting List Registrations 07/01/2019 to 06/30/2020 (%)			All Waiting List Registrations on 06/30/2020 (%)		
	This Center (N=306)	OPTN Region (N=2,751)	U.S. (N=39,776)	This Center (N=504)	OPTN Region (N=7,678)	U.S. (N=99,301)
All (%)	100.0	100.0	100.0	100.0	100.0	100.0
Blood Type (%)						
O	41.8	47.7	49.1	48.4	51.3	53.8
A	30.1	30.8	32.4	27.8	27.4	27.2
B	21.2	17.3	14.8	18.8	17.9	16.5
AB	6.9	4.2	3.8	5.0	3.4	2.5
Unknown	0.0	0.0	0.0	0.0	0.0	0.0
Previous Transplant (%)						
Yes	11.8	13.9	12.7	15.1	15.0	13.7
No	88.2	86.1	87.3	84.9	85.0	86.3
Unknown	0.0	0.0	0.0	0.0	0.0	0.0
Initial CPRA (%)						
0-9%	94.8	88.4	80.5	95.8	88.8	80.0
10-79%	2.9	7.3	12.4	1.8	7.0	12.6
80+%	2.3	4.3	6.9	2.4	4.2	7.3
Unknown	0.0	0.0	0.2	0.0	0.0	0.1
Primary Disease (%)*						
Glomerular Diseases	22.9	18.1	18.8	25.4	17.5	18.8
Tubular and Interstitial Diseases	5.2	4.2	3.8	5.8	3.8	3.6
Polycystic Kidneys	6.2	7.3	7.5	6.9	6.4	6.9
Congenital, Familial, Metabolic	2.0	1.9	2.1	1.8	1.5	1.8
Diabetes	37.3	34.1	35.4	32.9	36.0	36.9
Renovascular & Vascular Diseases	0.3	0.1	0.2	0.2	0.1	0.2
Neoplasms	0.3	0.5	0.3	0.2	0.4	0.3
Hypertensive Nephrosclerosis	16.3	21.4	19.8	16.3	23.9	21.2
Other	9.5	11.8	11.7	10.5	10.0	9.9
Missing*	0.0	0.5	0.4	0.0	0.4	0.4

* When "retransplant" is indicated, the primary disease is passed forward from the prior transplant in order to indicate the initial primary disease causing organ failure. "Missing" may include some patients for whom retransplant is indicated but no prior diagnosis can be found.



B. Waiting List Information

Table B4. Transplant rates: 07/01/2018 - 03/12/2020

Waiting List Registrations	This Center	OPO/DSA	Region	U.S.
All Candidates				
Count on waiting list at start*	344	6,508	7,650	100,697
Person Years**	737.0	11,260.3	13,213.1	171,846.3
Removals for Transplant	347	2,252	2,877	38,985
Adult (18+) Candidates				
Count on waiting list at start*	344	6,370	7,497	99,206
Person Years**	733.4	11,031.1	12,954.7	169,185.1
Removals for transplant	342	2,163	2,781	37,536
Pediatric (<18) Candidates				
Count on waiting list at start*	0	138	153	1,491
Person Years**	3.6	229.3	258.5	2,661.2
Removals for transplant	5	89	96	1,449

* Counts in this table may be lower than similar counts in other waiting list tables, such as Table B1. A small percentage (~1%) of patients are found to have died or been transplanted before being removed from the waiting list, so these patients are excluded if the event occurs prior to the start of the study period. Inactive time on the waiting list is included in the calculations for this table.

** Person years are calculated as days (converted to fractional years). The number of days from July 1 or from the date of first wait listing until death, transplant, removal from the waiting list or March 12, 2020.

Figure B1. Observed and expected transplant rates: 07/01/2018 - 03/12/2020




Figure B2. Transplant rate ratio estimate





Figure B3. Observed adult (18+) and pediatric (<18) transplant rates: 07/01/2018 - 03/12/2020





B. Waiting List Information

Table B4D. Deceased donor transplant rates: 07/01/2018 - 03/12/2020

Waiting List Registrations	This Center	OPO/DSA	Region	U.S.
All Candidates				
Count on waiting list at start*	344	6,508	7,650	100,697
Person Years**	737.0	11,260.3	13,213.1	171,846.3
Removals for Transplant	247	1,319	1,822	27,603
Adult (18+) Candidates				
Count on waiting list at start*	344	6,370	7,497	99,206
Person Years**	733.4	11,031.1	12,954.7	169,185.1
Removals for transplant	245	1,262	1,762	26,609
Pediatric (<18) Candidates				
Count on waiting list at start*	0	138	153	1,491
Person Years**	3.6	229.3	258.5	2,661.2
Removals for transplant	2	57	60	994

* Counts in this table may be lower than similar counts in other waiting list tables, such as Table B1. A small percentage (~1%) of patients are found to have died or been transplanted before being removed from the waiting list, so these patients are excluded if the event occurs prior to the start of the study period. Inactive time on the waiting list is included in the calculations for this table.

** Person years are calculated as days (converted to fractional years). The number of days from July 1 or from the date of first wait listing until death, transplant, removal from the waiting list or March 12, 2020.

Figure B1D. Observed and expected deceased donor transplant rates: 07/01/2018 - 03/12/2020




Figure B2D. Deceased donor transplant rate ratio estimate





Figure B3D. Observed adult (18+) and pediatric (<18) deceased donor transplant rates: 07/01/2018 - 03/12/2020





B. Waiting List Information

Table B5. Waiting list mortality rates: 07/01/2018 - 03/12/2020

Waiting List Registrations	This Center	OPO/DSA	Region	U.S.
All Candidates				
Count on waiting list at start*	344	6,508	7,650	100,697
Person Years**	774.6	11,834.8	13,965.9	184,662.2
Number of deaths	33	545	641	8,721
Adult (18+) Candidates				
Count on waiting list at start*	344	6,370	7,497	99,206
Person Years**	771.0	11,599.8	13,700.3	181,914.0
Number of deaths	33	540	635	8,686
Pediatric (<18) Candidates				
Count on waiting list at start*	0	138	153	1,491
Person Years**	3.6	235.0	265.6	2,748.2
Number of deaths	0	5	6	35

* Counts in this table may be lower than similar counts in other waiting list tables, such as Table B1. A small percentage (~1%) of patients are found to have died or been transplanted before being removed from the waiting list, so these patients are excluded if the event occurs prior to the start of the study period. Inactive time on the waiting list is included in the calculations for this table.

** Person years are calculated as days (converted to fractional years). The number of days from July 1 or from the date of first wait listing until death, transplant, 60 days after recovery, transfer or March 12, 2020.

Figure B4. Observed and expected waiting list mortality rates: 07/01/2018 - 03/12/2020




Figure B5. Waiting list mortality rate ratio estimate





Figure B6. Observed adult (18+) and pediatric (<18) waiting list mortality rates: 07/01/2018 - 03/12/2020





B. Waiting List Information

Table B6. Rates of patient mortality after listing: 07/01/2018 - 03/12/2020

Waiting List Registrations	This Center	OPO/DSA	Region	U.S.
All Patients				
Count at risk during the evaluation period	1,364	16,924	20,902	288,816
Person-years*	1,691.3	22,175.8	27,348.9	381,306.4
Number of Deaths	53	902	1,121	16,073
Adult (18+) Patients				
Count at risk during the evaluation period	1,353	16,422	20,305	280,336
Person-years*	1,680.6	21,495.9	26,543.0	369,696.7
Number of Deaths	53	895	1,113	16,004
Pediatric (<18) Patients				
Count at risk during the evaluation period	11	502	597	8,480
Person-years*	10.6	680.0	805.9	11,609.7
Number of Deaths	0	7	8	69

* Person-years are calculated as days (converted to fractional years). The number of days from 07/01/2018, or from the date of first wait listing until death, reaching 7 years after listing or March 12, 2020.

** Patient mortality after listing describes the relative survival experience of patients after listing. It depends on many factors, some of which are outside of the control of the transplant program. For example, availability of organs may not be the same in every part of the country.

Figure B7. Observed and expected rates of patient mortality after listing: 07/01/2018 - 03/12/2020




Figure B8. HR estimate of patient mortality after listing





Figure B9. Observed adult (18+) and pediatric (<18) rates of patient mortality after listing: 07/01/2018 - 03/12/2020





B. Waiting List Information

Table B7. Waiting list candidate status after listing
Candidates registered on waiting list between 01/01/2018 and 12/31/2018

Waiting list status (survival status)	This Center (N=333)			U.S. (N=39,987)		
	Months Since Listing			Months Since Listing		
	6	12	18	6	12	18
Alive on waiting list (%)	60.1	46.8	40.2	77.4	64.2	54.1
Died on the waiting list without transplant (%)	2.1	2.7	3.3	1.0	1.9	2.8
Removed without transplant (%):						
Condition worsened (status unknown)	0.9	1.2	2.4	0.6	1.4	2.4
Condition improved (status unknown)	0.0	0.0	0.0	0.1	0.1	0.2
Refused transplant (status unknown)	0.3	0.3	0.3	0.0	0.1	0.2
Other	1.2	2.4	2.7	0.8	1.6	2.6
Transplant (living donor from waiting list only) (%):						
Functioning (alive)	11.7	16.2	11.1	6.7	10.1	7.7
Failed-Retransplanted (alive)	0.0	0.0	0.0	0.0	0.0	0.0
Failed-alive not retransplanted	0.0	0.0	0.0	0.0	0.0	0.0
Died	0.0	0.6	0.9	0.0	0.1	0.1
Status Yet Unknown**	0.3	0.3	6.3	0.1	0.6	5.0
Transplant (deceased donor) (%):						
Functioning (alive)	21.3	24.3	18.9	11.1	14.6	11.6
Failed-Retransplanted (alive)	0.0	0.0	0.0	0.0	0.0	0.0
Failed-alive not retransplanted	0.0	0.0	0.0	0.0	0.0	0.1
Died	0.0	0.0	0.0	0.2	0.4	0.7
Status Yet Unknown*	1.5	4.5	13.2	1.7	4.0	11.4
Lost or Transferred (status unknown) (%)	0.6	0.6	0.6	0.2	0.7	1.3
TOTAL (%)	100.0	100.0	100.0	100.0	100.0	100.0
Total % known died on waiting list or after transplant	2.1	3.3	4.2	1.2	2.4	3.6
Total % known died or removed as unstable	3.0	4.5	6.6	1.9	3.8	6.0
Total % removed for transplant	34.8	45.9	50.5	19.9	29.9	36.5
Total % with known functioning transplant (alive)	33.0	40.5	30.0	17.8	24.7	19.3

* Follow-up form covering specified time period not yet completed, and possibly has not become due.



B. Waiting List Information

Table B8. Percent of candidates with deceased donor transplants: demographic characteristics
Candidates registered on the waiting list between 07/01/2014 and 06/30/2017

Characteristic	Percent transplanted at time periods since listing									
	This Center					United States				
	N	30 day	1 year	2 years	3 years	N	30 day	1 year	2 years	3 years
All	297	4.7	21.2	34.0	43.1	93,662	3.6	16.0	22.6	27.8
Ethnicity/Race*										
White	48	10.4	25.0	37.5	41.7	37,390	3.9	17.0	23.7	28.8
African-American	92	1.1	19.6	38.0	46.7	29,527	3.5	16.2	22.7	27.9
Hispanic/Latino	62	4.8	22.6	33.9	45.2	17,534	3.7	15.6	22.0	27.4
Asian	94	5.3	20.2	28.7	39.4	7,645	2.0	11.2	17.6	23.0
Other	1	0.0	0.0	0.0	0.0	1,566	4.5	19.5	25.9	31.6
Unknown	0	--	--	--	--	0	--	--	--	--
Age										
<2 years	0	--	--	--	--	128	5.5	36.7	53.9	68.0
2-11 years	0	--	--	--	--	852	8.3	48.4	62.0	70.8
12-17 years	0	--	--	--	--	1,407	7.8	49.3	62.2	67.9
18-34 years	32	3.1	21.9	34.4	40.6	9,563	3.5	17.6	25.9	32.9
35-49 years	73	6.8	17.8	30.1	45.2	23,762	3.3	15.0	21.9	27.7
50-64 years	146	3.4	21.2	34.9	43.8	40,111	3.6	14.8	20.7	25.5
65-69 years	37	5.4	29.7	43.2	45.9	12,262	3.7	14.8	20.6	25.1
70+ years	9	11.1	11.1	11.1	11.1	5,577	3.3	15.3	20.9	25.1
Gender										
Male	192	3.6	18.2	32.8	42.7	58,067	3.8	15.6	21.7	26.8
Female	105	6.7	26.7	36.2	43.8	35,595	3.4	16.8	24.0	29.5

* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



B. Waiting List Information

Table B9. Percent of candidates with deceased donor transplants: medical characteristics
Candidates registered on the waiting list between 07/01/2014 and 06/30/2017

Characteristic	N	Percent transplanted at time periods since listing				N	United States			
		This Center					30 day	1 year	2 years	3 years
		30 day	1 year	2 years	3 years					
All	297	4.7	21.2	34.0	43.1	93,662	3.6	16.0	22.6	27.8
Blood Type										
O	139	5.0	18.7	33.1	43.2	46,625	3.3	14.0	19.5	24.0
A	84	4.8	17.9	31.0	42.9	29,375	4.4	19.0	26.9	33.2
B	60	5.0	28.3	35.0	38.3	14,175	2.4	13.1	19.2	24.0
AB	14	0.0	35.7	57.1	64.3	3,487	6.0	30.6	41.5	49.3
Previous Transplant										
Yes	28	0.0	14.3	17.9	25.0	12,856	2.7	16.4	24.4	29.8
No	269	5.2	21.9	35.7	45.0	80,806	3.8	16.0	22.3	27.5
Peak PRA/CPRA										
0-9%	290	4.8	20.7	33.8	43.1	75,317	3.9	15.5	21.6	26.8
10-79%	3	0.0	33.3	33.3	33.3	10,787	2.5	15.7	22.8	28.2
80+%	4	0.0	50.0	50.0	50.0	7,475	2.6	22.4	32.3	37.4
Unknown	0	--	--	--	--	7	100.0	100.0	100.0	100.0
Primary Disease*										
Glomerular Diseases	82	0.0	14.6	22.0	42.7	17,106	2.9	16.8	25.0	31.6
Tubular & Interstitial Diseases	13	0.0	7.7	23.1	23.1	3,474	4.6	19.5	27.3	32.6
Polycystic Kidneys	22	0.0	22.7	40.9	45.5	6,196	2.1	14.6	22.1	28.9
Congenital, Familial, Metabolic	3	0.0	33.3	33.3	33.3	1,840	4.9	29.2	39.0	46.6
Diabetes	91	2.2	22.0	35.2	41.8	32,973	2.4	11.7	16.6	20.7
Renovascular & Vascular Diseases	0	--	--	--	--	142	6.3	20.4	27.5	33.8
Neoplasms	0	--	--	--	--	288	5.6	22.2	28.5	34.7
Hypertensive Nephrosclerosis	47	4.3	19.1	36.2	40.4	20,390	3.4	15.9	22.6	28.2
Other	39	25.6	38.5	53.8	56.4	10,889	9.1	25.7	32.9	37.5
Missing*	0	--	--	--	--	364	1.4	11.0	16.5	20.9

* When "retransplant" is indicated, the primary disease is passed forward from the prior transplant in order to indicate the initial primary disease causing organ failure. "Missing" may include some patients for whom retransplant is indicated but no prior diagnosis can be found.



B. Waiting List Information

Table B10. Time to transplant for waiting list candidates*
Candidates registered on the waiting list between 07/01/2014 and 12/31/2019

Percentile	Center	Months to Transplant**		U.S.
		OPO/DSA	Region	
5th	0.4	0.3	0.4	0.9
10th	1.3	1.4	1.5	2.5
25th	4.7	9.0	8.5	9.7
50th (median time to transplant)	20.1	55.1	45.0	42.4
75th	Not Observed	Not Observed	Not Observed	Not Observed

* If cells contain "Not Observed" fewer than that percentile of patients had received a transplant. For example, the 50th percentile of time to transplant is the time when 50% of candidates have received transplants. If waiting times are long, then the 50th percentile may not be observed during the follow-up period for this table. Also, if more than 50% of candidates are removed from the list due to death or other reasons before receiving transplants, then the 50th percentile of time to transplant will not be observed.

** Censored on 06/30/2020. Calculated as the months after listing, during which the corresponding percent of all patients initially listed had received a transplant.



B. Waiting List Information

Table B11. Offer Acceptance Practices: 07/01/2019 - 03/12/2020

Offers Acceptance Characteristics	This Center	OPO/DSA	Region	U.S.
Overall				
Number of Offers	7,611	124,053	132,631	1,489,166
Number of Acceptances	97	473	691	11,332
Expected Acceptances	20.1	441.2	561.8	11,315.1
Offer Acceptance Ratio*	4.48	1.07	1.23	1.00
95% Credible Interval**	[3.64, 5.41]	--	--	--
Low-KDRI Donors (KDRI < 1.05)				
Number of Offers	1,064	14,977	16,083	148,107
Number of Acceptances	17	102	164	3,735
Expected Acceptances	4.6	100.2	145.5	3,727.8
Offer Acceptance Ratio*	2.86	1.02	1.13	1.00
95% Credible Interval**	[1.72, 4.29]	--	--	--
Medium-KDRI Donors (1.05 < KDRI < 1.75)				
Number of Offers	4,234	81,671	86,896	1,024,741
Number of Acceptances	64	285	423	6,217
Expected Acceptances	10.6	249.6	310.5	6,209.0
Offer Acceptance Ratio*	5.24	1.14	1.36	1.00
95% Credible Interval**	[4.05, 6.57]	--	--	--
High-KDRI Donors (KDRI > 1.75)				
Number of Offers	2,313	27,405	29,652	316,318
Number of Acceptances	16	86	104	1,380
Expected Acceptances	4.8	91.3	105.7	1,378.3
Offer Acceptance Ratio*	2.63	0.94	0.98	1.00
95% Credible Interval**	[1.56, 3.97]	--	--	--
Hard-to-Place Kidneys (Over 100 Offers)				
Number of Offers	7,276	112,783	119,656	1,289,977
Number of Acceptances	72	210	268	1,603
Expected Acceptances	8.1	140.7	146.4	1,598.9
Offer Acceptance Ratio*	7.34	1.49	1.82	1.00
95% Credible Interval**	[5.77, 9.11]	--	--	--

* The offer acceptance ratio estimates the relative offer acceptance practice of New York University Medical Center (NYUC) compared to the national offer acceptance practice. A ratio above one indicates the program is more likely to accept an offer compared to national offer acceptance practices (e.g., an offer acceptance ratio of 1.25 indicates a 25% more likely to accept an offer), while a ratio below one indicates the program is less likely to accept an offer compared to national offer acceptance practices (e.g., an offer acceptance ratio of 0.75 indicates a 25% less likely to accept an offer).

** As an example, the 95% Credible Interval for the overall offer acceptance ratio, [3.64, 5.41], indicates the location of NYUC's true offer acceptance ratio with 95% probability. The best estimate is 348% more likely to accept an offer compared to national acceptance behavior, but NYUC's performance could plausibly range from 264% higher acceptance up to 441% higher acceptance.



B. Waiting List Information

Figure B10. Offer acceptance: Overall




Figure B11. Offer acceptance: Low-KDRI




Figure B12. Offer acceptance: Medium-KDRI




Figure B13. Offer acceptance: High-KDRI





Figure B14. Offer acceptance: Offer number > 100





C. Transplant Information

Table C1D. Deceased donor transplant recipient demographic characteristics
Patients transplanted between 07/01/2019 and 06/30/2020

Characteristic	Percentage in each category		
	Center (N=136)	Region (N=953)	U.S. (N=16,870)
Ethnicity/Race (%)*			
White	25.7	37.1	37.8
African-American	35.3	33.7	32.2
Hispanic/Latino	17.6	17.8	20.3
Asian	20.6	10.3	7.8
Other	0.7	1.0	1.9
Unknown	0.0	0.0	0.0
Age (%)			
<2 years	0.0	0.0	0.1
2-11 years	0.0	0.8	1.2
12-17	0.0	0.8	1.6
18-34	8.8	9.3	10.3
35-49 years	24.3	21.7	23.9
50-64 years	37.5	43.4	40.3
65-69 years	16.2	14.3	13.2
70+ years	13.2	9.5	9.5
Gender (%)			
Male	72.8	64.4	60.4
Female	27.2	35.6	39.6

* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



C. Transplant Information

Table C1L. Living donor transplant recipient demographic characteristics
Patients transplanted between 07/01/2019 and 06/30/2020

Characteristic	Percentage in each category		
	Center (N=50)	Region (N=501)	U.S. (N=5,777)
Ethnicity/Race (%)*			
White	62.0	63.3	65.1
African-American	14.0	13.6	12.4
Hispanic/Latino	10.0	15.8	14.7
Asian	14.0	7.0	6.4
Other	0.0	0.4	1.4
Unknown	0.0	0.0	0.0
Age (%)			
<2 years	0.0	0.0	0.2
2-11 years	0.0	0.6	1.7
12-17	0.0	2.0	1.7
18-34	12.0	15.0	15.9
35-49 years	18.0	23.8	25.7
50-64 years	26.0	35.3	35.3
65-69 years	24.0	12.4	10.9
70+ years	20.0	11.0	8.7
Gender (%)			
Male	64.0	65.1	62.3
Female	36.0	34.9	37.7

* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



C. Transplant Information

Table C2D. Deceased donor transplant recipient medical characteristics
Patients transplanted between 07/01/2019 and 06/30/2020

Characteristic	Percentage in each category		
	Center (N=136)	Region (N=953)	U.S. (N=16,870)
Blood Type (%)			
O	44.9	46.2	45.5
A	30.1	34.1	35.2
B	20.6	14.9	14.1
AB	4.4	4.8	5.3
Previous Transplant (%)			
Yes	7.4	14.5	12.6
No	92.6	85.5	87.4
Peak PRA/CPRA Prior to Transplant (%)			
0-9%	86.8	69.3	61.2
10-79%	5.9	16.8	22.8
80+ %	7.4	14.0	16.0
Unknown	0.0	0.0	0.0
Body Mass Index (%)			
0-20	10.3	10.1	9.4
21-25	36.8	31.8	27.1
26-30	29.4	29.5	30.6
31-35	14.7	17.6	21.1
36-40	4.4	7.1	8.1
41+	0.0	2.3	1.5
Unknown	4.4	1.6	2.2
Primary Disease (%)*			
Glomerular Diseases	25.7	21.9	21.2
Tubular and Interstitial Disease	3.7	3.9	4.0
Polycystic Kidneys	5.1	7.9	7.5
Congenital, Familial, Metabolic	0.0	1.4	2.6
Diabetes	35.3	28.5	30.1
Renovascular & Vascular Diseases	0.0	0.2	0.2
Neoplasms	0.0	0.4	0.4
Hypertensive Nephrosclerosis	19.1	21.9	23.0
Other Kidney	11.0	13.6	10.7
Missing*	0.0	0.2	0.2

* When "retransplant" is indicated, the primary disease is passed forward from the prior transplant in order to indicate the initial primary disease causing organ failure. "Missing" may include some patients for whom retransplant is indicated but no prior diagnosis can be found.



C. Transplant Information

Table C2L. Living donor transplant recipient medical characteristics
Patients transplanted between 07/01/2019 and 06/30/2020

Characteristic	Percentage in each category		
	Center (N=50)	Region (N=501)	U.S. (N=5,777)
Blood Type (%)			
O	36.0	42.3	43.4
A	34.0	34.9	38.8
B	24.0	18.8	14.1
AB	6.0	4.0	3.7
Previous Transplant (%)			
Yes	12.0	12.4	10.3
No	88.0	87.6	89.7
Peak PRA/CPRA Prior to Transplant (%)			
0-9%	84.0	83.6	74.8
10-79%	14.0	13.2	20.0
80+ %	2.0	3.2	5.2
Unknown	0.0	0.0	0.0
Body Mass Index (%)			
0-20	12.0	11.0	11.3
21-25	28.0	28.3	27.7
26-30	38.0	31.1	31.6
31-35	16.0	21.4	19.5
36-40	6.0	6.0	7.7
41+	0.0	1.0	1.1
Unknown	0.0	1.2	1.1
Primary Disease (%)*			
Glomerular Diseases	24.0	27.7	28.0
Tubular and Interstitial Disease	8.0	6.2	5.5
Polycystic Kidneys	10.0	10.0	13.1
Congenital, Familial, Metabolic	2.0	2.8	3.8
Diabetes	34.0	25.0	24.8
Renovascular & Vascular Diseases	0.0	0.0	0.3
Neoplasms	0.0	0.2	0.4
Hypertensive Nephrosclerosis	12.0	17.0	15.1
Other Kidney	10.0	10.8	8.9
Missing*	0.0	0.4	0.2

* When "retransplant" is indicated, the primary disease is passed forward from the prior transplant in order to indicate the initial primary disease causing organ failure. "Missing" may include some patients for whom retransplant is indicated but no prior diagnosis can be found.



C. Transplant Information

Table C3D. Deceased donor characteristics
Transplants performed between 07/01/2019 and 06/30/2020

Donor Characteristic	Percentage in each category		
	Center (N=136)	Region (N=953)	U.S. (N=16,870)
Cause of Death (%)			
Deceased: Stroke	19.1	18.9	22.7
Deceased: MVA	7.4	9.0	13.3
Deceased: Other	73.5	72.1	63.9
Ethnicity/Race (%)*			
White	66.9	68.0	66.8
African-American	14.0	12.9	13.2
Hispanic/Latino	13.2	14.4	15.8
Asian	4.4	4.1	2.9
Other	1.5	0.6	1.2
Not Reported	0.0	0.0	0.0
Age (%)			
<2 years	1.5	1.0	0.8
2-11 years	1.5	2.6	2.5
12-17	0.7	2.9	3.8
18-34	35.3	31.3	33.6
35-49 years	34.6	31.4	31.7
50-64 years	25.0	27.8	25.2
65-69 years	1.5	2.4	2.0
70+ years	0.0	0.5	0.5
Gender (%)			
Male	63.2	62.1	63.0
Female	36.8	37.9	37.0
Blood Type (%)			
O	48.5	48.8	47.2
A	39.0	36.2	37.6
B	11.8	11.5	11.7
AB	0.7	3.5	3.5
Unknown	0.0	0.0	0.0

* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



C. Transplant Information

Table C3L. Living donor characteristics
Transplants performed between 07/01/2019 and 06/30/2020

Donor Characteristic	Percentage in each category		
	Center (N=50)	Region (N=501)	U.S. (N=5,777)
Ethnicity/Race (%)*			
White	66.0	65.1	71.5
African-American	14.0	13.8	7.9
Hispanic/Latino	8.0	13.6	14.0
Asian	10.0	7.2	4.9
Other	2.0	0.4	1.6
Not Reported	0.0	0.0	0.0
Age (%)			
0-11 years	0.0	0.0	0.0
12-17	0.0	0.0	0.0
18-34	42.0	26.3	25.9
35-49 years	42.0	39.5	38.3
50-64 years	12.0	28.7	29.8
65-69 years	0.0	3.6	4.5
70+ years	4.0	1.8	1.5
Gender (%)			
Male	48.0	40.7	35.1
Female	52.0	59.3	64.9
Blood Type (%)			
O	48.0	62.5	62.6
A	26.0	23.4	27.4
B	20.0	12.8	8.6
AB	6.0	1.4	1.4
Unknown	0.0	0.0	0.0

* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



C. Transplant Information

Table C4D. Deceased donor transplant characteristics
Transplants performed between 07/01/2019 and 06/30/2020

Transplant Characteristic	Percentage in each category		
	Center (N=136)	Region (N=953)	U.S. (N=16,870)
Cold Ischemic Time (Hours): Local (%)			
Deceased: 0-11 hr	17.1	34.1	34.3
Deceased: 12-21 hr	40.0	48.3	47.8
Deceased: 22-31 hr	40.0	15.3	14.6
Deceased: 32-41 hr	2.9	1.1	1.6
Deceased: 42+ hr	0.0	0.4	0.5
Not Reported	0.0	0.8	1.2
Cold Ischemic Time (Hours): Shared (%)			
Deceased: 0-11 hr	5.9	10.2	9.2
Deceased: 12-21 hr	4.0	15.4	36.3
Deceased: 22-31 hr	48.5	37.8	38.4
Deceased: 32-41 hr	34.7	26.6	11.7
Deceased: 42+ hr	5.0	6.9	3.1
Not Reported	2.0	3.1	1.3
Level of Mismatch (%)			
A Locus Mismatches (%)			
0	5.9	11.1	11.2
1	39.0	36.8	39.7
2	55.1	51.5	48.9
Not Reported	0.0	0.5	0.2
B Locus Mismatches (%)			
0	2.9	6.7	7.0
1	19.9	23.1	25.0
2	77.2	69.7	67.7
Not Reported	0.0	0.5	0.2
DR Locus Mismatches (%)			
0	9.6	13.0	16.0
1	41.9	45.1	47.7
2	48.5	41.3	36.1
Not Reported	0.0	0.5	0.2
Total Mismatches (%)			
0	1.5	4.8	4.5
1	1.5	1.3	1.1
2	2.2	3.4	4.7
3	8.8	11.8	13.9
4	22.8	22.9	27.3
5	40.4	37.0	33.0
6	22.8	18.4	15.2
Not Reported	0.0	0.5	0.2
Procedure Type (%)			
Kidney alone	92.6	94.1	93.8
Kidney and another organ	7.4	5.9	6.2
Dialysis in First Week After Transplant (%)			
Yes	46.3	34.7	28.6
No	53.7	65.0	70.9
Not Reported	0.0	0.3	0.5
Sharing (%)			
Local	25.7	49.5	69.6
Shared	74.3	50.5	30.4
Median Time in Hospital After Transplant*	7.0 Days	6.0 Days	5.0 Days

* Multiple organ transplants are excluded from this statistic.



C. Transplant Information

Table C4L. Living donor transplant characteristics
Transplants performed between 07/01/2019 and 06/30/2020

Transplant Characteristic	Percentage in each category		
	Center (N=50)	Region (N=501)	U.S. (N=5,777)
Relation with Donor (%)			
Related	46.0	42.5	37.9
Unrelated	54.0	57.5	61.7
Not Reported	0.0	0.0	0.3
Level of Mismatch (%)			
A Locus Mismatches (%)			
0	20.0	15.0	16.2
1	54.0	50.5	48.6
2	24.0	33.7	31.6
Not Reported	2.0	0.8	3.5
B Locus Mismatches (%)			
0	4.0	10.0	9.7
1	62.0	47.3	41.6
2	32.0	41.9	45.1
Not Reported	2.0	0.8	3.6
DR Locus Mismatches (%)			
0	10.0	18.2	15.1
1	64.0	52.3	48.4
2	24.0	28.7	33.0
Not Reported	2.0	0.8	3.5
Total Mismatches (%)			
0	2.0	4.2	4.3
1	4.0	4.6	4.1
2	14.0	13.0	11.5
3	40.0	27.3	23.0
4	14.0	14.4	18.2
5	12.0	25.3	22.8
6	12.0	10.4	12.5
Not Reported	2.0	0.8	3.6
Procedure Type (%)			
Kidney alone	100.0	100.0	100.0
Kidney and another organ	0.0	0.0	0.0
Dialysis in First Week After Transplant (%)			
Yes	2.0	5.0	2.7
No	98.0	95.0	96.8
Not Reported	0.0	0.0	0.5
Median Time in Hospital After Transplant*	5.0 Days	4.0 Days	4.0 Days

* Multiple organ transplants are excluded from this statistic.



C. Transplant Information

Table C5. Adult (18+) 1-month survival with a functioning graft
Single organ transplants performed between 07/01/2017 and 12/31/2019
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	420	50,251
Estimated probability of surviving with a functioning graft at 1 month (unadjusted for patient and donor characteristics)	99.05%	98.67%
Expected probability of surviving with a functioning graft at 1 month (adjusted for patient and donor characteristics)	98.41%	--
Number of observed graft failures (including deaths) during the first month after transplant	4	669
Number of expected graft failures (including deaths) during the first month after transplant	6.70	--
Estimated hazard ratio*	0.69	--
95% credible interval for the hazard ratio**	[0.25, 1.34]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.25, 1.34], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 31% lower risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 75% reduced risk up to 34% increased risk.

Figure C1. Adult (18+) 1-month graft failure HR estimate





Figure C2. Adult (18+) 1-month graft failure HR program comparison





C. Transplant Information

Table C5D. Adult (18+) 1-month survival with a functioning deceased donor graft
Single organ transplants performed between 07/01/2017 and 12/31/2019
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	295	34,632
Estimated probability of surviving with a functioning graft at 1 month (unadjusted for patient and donor characteristics)	98.98%	98.42%
Expected probability of surviving with a functioning graft at 1 month (adjusted for patient and donor characteristics)	98.07%	--
Number of observed graft failures (including deaths) during the first month after transplant	3	547
Number of expected graft failures (including deaths) during the first month after transplant	5.73	--
Estimated hazard ratio*	0.65	--
95% credible interval for the hazard ratio**	[0.21, 1.33]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.21, 1.33], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 35% lower risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 79% reduced risk up to 33% increased risk.

Figure C1D. Adult (18+) 1-month deceased donor graft failure HR estimate





Figure C2D. Adult (18+) 1-month deceased donor graft failure HR program comparison





C. Transplant Information

Table C5L. Adult (18+) 1-month survival with a functioning living donor graft
Single organ transplants performed between 07/01/2017 and 12/31/2019
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	125	15,619
Estimated probability of surviving with a functioning graft at 1 month (unadjusted for patient and donor characteristics)	99.20%	99.22%
Expected probability of surviving with a functioning graft at 1 month (adjusted for patient and donor characteristics)	99.23%	--
Number of observed graft failures (including deaths) during the first month after transplant	1	122
Number of expected graft failures (including deaths) during the first month after transplant	0.97	--
Estimated hazard ratio*	1.01	--
95% credible interval for the hazard ratio**	[0.21, 2.43]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.21, 2.43], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 1% higher risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 79% reduced risk up to 143% increased risk.

Figure C1L. Adult (18+) 1-month living donor graft failure HR estimate





Figure C2L. Adult (18+) 1-month living donor graft failure HR program comparison





C. Transplant Information

Table C6. Adult (18+) 1-year survival with a functioning graft
Single organ transplants performed between 07/01/2017 and 12/31/2019
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	420	50,251
Estimated probability of surviving with a functioning graft at 1 year (unadjusted for patient and donor characteristics)	96.50%	95.75%
Expected probability of surviving with a functioning graft at 1 year (adjusted for patient and donor characteristics)	94.89%	--
Number of observed graft failures (including deaths) during the first year after transplant	14	1,946
Number of expected graft failures (including deaths) during the first year after transplant	19.78	--
Estimated hazard ratio*	0.73	--
95% credible interval for the hazard ratio**	[0.42, 1.14]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.42, 1.14], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 27% lower risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 58% reduced risk up to 14% increased risk.

Figure C3. Adult (18+) 1-year graft failure HR estimate





Figure C4. Adult (18+) 1-year graft failure HR program comparison





C. Transplant Information

Table C6D. Adult (18+) 1-year survival with a functioning deceased donor graft
Single organ transplants performed between 07/01/2017 and 12/31/2019
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	295	34,632
Estimated probability of surviving with a functioning graft at 1 year (unadjusted for patient and donor characteristics)	96.41%	94.73%
Expected probability of surviving with a functioning graft at 1 year (adjusted for patient and donor characteristics)	93.56%	--
Number of observed graft failures (including deaths) during the first year after transplant	10	1,661
Number of expected graft failures (including deaths) during the first year after transplant	17.57	--
Estimated hazard ratio*	0.61	--
95% credible interval for the hazard ratio**	[0.32, 1.01]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.32, 1.01], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 39% lower risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 68% reduced risk up to 1% increased risk.

Figure C3D. Adult (18+) 1-year deceased donor graft failure HR estimate





Figure C4D. Adult (18+) 1-year deceased donor graft failure HR program comparison





C. Transplant Information

Table C6L. Adult (18+) 1-year survival with a functioning living donor graft
Single organ transplants performed between 07/01/2017 and 12/31/2019
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	125	15,619
Estimated probability of surviving with a functioning graft at 1 year (unadjusted for patient and donor characteristics)	96.74%	98.02%
Expected probability of surviving with a functioning graft at 1 year (adjusted for patient and donor characteristics)	98.03%	--
Number of observed graft failures (including deaths) during the first year after transplant	4	285
Number of expected graft failures (including deaths) during the first year after transplant	2.21	--
Estimated hazard ratio*	1.42	--
95% credible interval for the hazard ratio**	[0.52, 2.77]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.52, 2.77], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 42% higher risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 48% reduced risk up to 177% increased risk.

Figure C3L. Adult (18+) 1-year living donor graft failure HR estimate





Figure C4L. Adult (18+) 1-year living donor graft failure HR program comparison





C. Transplant Information

Table C7. Adult (18+) 3-year survival with a functioning graft
Single organ transplants performed between 01/01/2015 and 06/30/2017
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	204	42,472
Estimated probability of surviving with a functioning graft at 3 years (unadjusted for patient and donor characteristics)	85.15%	89.69%
Expected probability of surviving with a functioning graft at 3 years (adjusted for patient and donor characteristics)	84.93%	--
Number of observed graft failures (including deaths) during the first 3 years after transplant	30	4,352
Number of expected graft failures (including deaths) during the first 3 years after transplant	30.18	--
Estimated hazard ratio*	0.99	--
95% credible interval for the hazard ratio**	[0.68, 1.37]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.68, 1.37], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 1% lower risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 32% reduced risk up to 37% increased risk.

Figure C5. Adult (18+) 3-year graft failure HR estimate





Figure C6. Adult (18+) 3-year graft failure HR program comparison





C. Transplant Information

Table C7D. Adult (18+) 3-year survival with a functioning deceased donor graft
Single organ transplants performed between 01/01/2015 and 06/30/2017
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	170	29,060
Estimated probability of surviving with a functioning graft at 3 years (unadjusted for patient and donor characteristics)	83.93%	87.51%
Expected probability of surviving with a functioning graft at 3 years (adjusted for patient and donor characteristics)	83.03%	--
Number of observed graft failures (including deaths) during the first 3 years after transplant	27	3,610
Number of expected graft failures (including deaths) during the first 3 years after transplant	28.30	--
Estimated hazard ratio*	0.96	--
95% credible interval for the hazard ratio**	[0.64, 1.34]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.64, 1.34], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 4% lower risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 36% reduced risk up to 34% increased risk.

Figure C5D. Adult (18+) 3-year deceased donor graft failure HR estimate





Figure C6D. Adult (18+) 3-year deceased donor graft failure HR program comparison





C. Transplant Information

Table C7L. Adult (18+) 3-year survival with a functioning living donor graft
Single organ transplants performed between 01/01/2015 and 06/30/2017
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	34	13,412
Estimated probability of surviving with a functioning graft at 3 years (unadjusted for patient and donor characteristics)	91.18%	94.42%
Expected probability of surviving with a functioning graft at 3 years (adjusted for patient and donor characteristics)	94.43%	--
Number of observed graft failures (including deaths) during the first 3 years after transplant	3	742
Number of expected graft failures (including deaths) during the first 3 years after transplant	1.88	--
Estimated hazard ratio*	1.29	--
95% credible interval for the hazard ratio**	[0.42, 2.64]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.42, 2.64], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 29% higher risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 58% reduced risk up to 164% increased risk.

Figure C5L. Adult (18+) 3-year living donor graft failure HR estimate





Figure C6L. Adult (18+) 3-year living donor graft failure HR program comparison





C. Transplant Information

Table C8. Pediatric (<18) 1-month survival with a functioning graft
Single organ transplants performed between 07/01/2017 and 12/31/2019
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	6	2,123
Estimated probability of surviving with a functioning graft at 1 month (unadjusted for patient and donor characteristics)	100.00%	98.92%
Expected probability of surviving with a functioning graft at 1 month (adjusted for patient and donor characteristics)	99.01%	--
Number of observed graft failures (including deaths) during the first month after transplant	0	23
Number of expected graft failures (including deaths) during the first month after transplant	0.06	--
Estimated hazard ratio*	0.97	--
95% credible interval for the hazard ratio**	[0.12, 2.71]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.12, 2.71], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 3% lower risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 88% reduced risk up to 171% increased risk.

Figure C7. Pediatric (<18) 1-month graft failure HR estimate





Figure C8. Pediatric (<18) 1-month graft failure HR program comparison





C. Transplant Information

Table C8D. Pediatric (<18) 1-month survival with a functioning deceased donor graft
Single organ transplants performed between 07/01/2017 and 12/31/2019
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	3	1,428
Estimated probability of surviving with a functioning graft at 1 month (unadjusted for patient and donor characteristics)	100.00%	98.74%
Expected probability of surviving with a functioning graft at 1 month (adjusted for patient and donor characteristics)	98.74%	--
Number of observed graft failures (including deaths) during the first month after transplant	0	18
Number of expected graft failures (including deaths) during the first month after transplant	0.04	--
Estimated hazard ratio*	0.98	--
95% credible interval for the hazard ratio**	[0.12, 2.73]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.12, 2.73], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 2% lower risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 88% reduced risk up to 173% increased risk.

Figure C7D. Pediatric (<18) 1-month deceased donor graft failure HR estimate





Figure C8D. Pediatric (<18) 1-month deceased donor graft failure HR program comparison





C. Transplant Information

Table C8L. Pediatric (<18) 1-month survival with a functioning living donor graft
Single organ transplants performed between 07/01/2017 and 12/31/2019
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	3	695
Estimated probability of surviving with a functioning graft at 1 month (unadjusted for patient and donor characteristics)	100.00%	99.28%
Expected probability of surviving with a functioning graft at 1 month (adjusted for patient and donor characteristics)	99.28%	--
Number of observed graft failures (including deaths) during the first month after transplant	0	5
Number of expected graft failures (including deaths) during the first month after transplant	0.02	--
Estimated hazard ratio*	0.99	--
95% credible interval for the hazard ratio**	[0.12, 2.76]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.12, 2.76], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 1% lower risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 88% reduced risk up to 176% increased risk.

Figure C7L. Pediatric (<18) 1-month living donor graft failure HR estimate





Figure C8L. Pediatric (<18) 1-month living donor graft failure HR program comparison





C. Transplant Information

Table C9. Pediatric (<18) 1-year survival with a functioning graft
Single organ transplants performed between 07/01/2017 and 12/31/2019
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	6	2,123
Estimated probability of surviving with a functioning graft at 1 year (unadjusted for patient and donor characteristics)	100.00%	98.18%
Expected probability of surviving with a functioning graft at 1 year (adjusted for patient and donor characteristics)	98.38%	--
Number of observed graft failures (including deaths) during the first year after transplant	0	36
Number of expected graft failures (including deaths) during the first year after transplant	0.10	--
Estimated hazard ratio*	0.95	--
95% credible interval for the hazard ratio**	[0.12, 2.66]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.12, 2.66], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 5% lower risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 88% reduced risk up to 166% increased risk.

Figure C9. Pediatric (<18) 1-year graft failure HR estimate





Figure C10. Pediatric (<18) 1-year graft failure HR program comparison





C. Transplant Information

Table C9D. Pediatric (<18) 1-year survival with a functioning deceased donor graft
Single organ transplants performed between 07/01/2017 and 12/31/2019
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	3	1,428
Estimated probability of surviving with a functioning graft at 1 year (unadjusted for patient and donor characteristics)	100.00%	97.80%
Expected probability of surviving with a functioning graft at 1 year (adjusted for patient and donor characteristics)	97.80%	--
Number of observed graft failures (including deaths) during the first year after transplant	0	29
Number of expected graft failures (including deaths) during the first year after transplant	0.06	--
Estimated hazard ratio*	0.97	--
95% credible interval for the hazard ratio**	[0.12, 2.70]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.12, 2.70], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 3% lower risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 88% reduced risk up to 170% increased risk.

Figure C9D. Pediatric (<18) 1-year deceased donor graft failure HR estimate





Figure C10D. Pediatric (<18) 1-year deceased donor graft failure HR program comparison





C. Transplant Information

Table C9L. Pediatric (<18) 1-year survival with a functioning living donor graft
Single organ transplants performed between 07/01/2017 and 12/31/2019
Deaths and retransplants are considered graft failures

	NYUC	U.S.
Number of transplants evaluated	3	695
Estimated probability of surviving with a functioning graft at 1 year (unadjusted for patient and donor characteristics)	100.00%	98.95%
Expected probability of surviving with a functioning graft at 1 year (adjusted for patient and donor characteristics)	98.95%	--
Number of observed graft failures (including deaths) during the first year after transplant	0	7
Number of expected graft failures (including deaths) during the first year after transplant	0.03	--
Estimated hazard ratio*	0.98	--
95% credible interval for the hazard ratio**	[0.12, 2.74]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.12, 2.74], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 2% lower risk of graft failure compared to an average program, but NYUC's performance could plausibly range from 88% reduced risk up to 174% increased risk.

Figure C9L. Pediatric (<18) 1-year living donor graft failure HR estimate





Figure C10L. Pediatric (<18) 1-year living donor graft failure HR program comparison





C. Transplant Information

Table C10. Pediatric (<18) 3-year survival with a functioning graft
Single organ transplants performed between 01/01/2015 and 06/30/2017
Deaths and retransplants are considered graft failures

This center did not perform any
transplants relevant to
this table during
01/01/2015-06/30/2017

**Figure C11. Pediatric (<18) 3-year
graft failure HR estimate**

This center did not perform any
transplants relevant to
this figure during
01/01/2015-06/30/2017

**Figure C12. Pediatric (<18) 3-year graft failure HR program
comparison**

This center did not perform any
transplants relevant to
this figure during
01/01/2015-06/30/2017



C. Transplant Information

Table C10D. Pediatric (<18) 3-year survival with a functioning deceased donor graft
Single organ transplants performed between 01/01/2015 and 06/30/2017
Deaths and retransplants are considered graft failures

This center did not perform any
transplants relevant to
this table during
01/01/2015-06/30/2017

Figure C11D. Pediatric (<18)
3-year deceased donor graft
failure HR estimate

This center did not perform any
transplants relevant to
this figure during
01/01/2015-06/30/2017

Figure C12D. Pediatric (<18) 3-year deceased donor graft
failure HR program comparison

This center did not perform any
transplants relevant to
this figure during
01/01/2015-06/30/2017



C. Transplant Information

Table C10L. Pediatric (<18) 3-year survival with a functioning living donor graft
Single organ transplants performed between 01/01/2015 and 06/30/2017
Deaths and retransplants are considered graft failures

This center did not perform any
transplants relevant to
this table during
01/01/2015-06/30/2017

Figure C11L. Pediatric (<18)
3-year living donor graft failure
HR estimate

This center did not perform any
transplants relevant to
this figure during
01/01/2015-06/30/2017

Figure C12L. Pediatric (<18) 3-year living donor graft failure
HR program comparison

This center did not perform any
transplants relevant to
this figure during
01/01/2015-06/30/2017



C. Transplant Information

Table C11. Adult (18+) 1-month patient survival
Single organ transplants performed between 07/01/2017 and 12/31/2019
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	389	44,658
Estimated probability of surviving at 1 month (unadjusted for patient and donor characteristics)	100.00%	99.56%
Expected probability of surviving at 1 month (adjusted for patient and donor characteristics)	99.51%	--
Number of observed deaths during the first month after transplant	0	198
Number of expected deaths during the first month after transplant	1.93	--
Estimated hazard ratio*	0.51	--
95% credible interval for the hazard ratio**	[0.06, 1.42]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.06, 1.42], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 49% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 94% reduced risk up to 42% increased risk.

Figure C13. Adult (18+) 1-month patient death HR estimate





Figure C14. Adult (18+) 1-month patient death HR program comparison





C. Transplant Information

Table C11D. Adult (18+) 1-month patient survival (deceased donor graft recipients)
Single organ transplants performed between 07/01/2017 and 12/31/2019
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	277	30,522
Estimated probability of surviving at 1 month (unadjusted for patient and donor characteristics)	100.00%	99.45%
Expected probability of surviving at 1 month (adjusted for patient and donor characteristics)	99.40%	--
Number of observed deaths during the first month after transplant	0	167
Number of expected deaths during the first month after transplant	1.67	--
Estimated hazard ratio*	0.54	--
95% credible interval for the hazard ratio**	[0.07, 1.52]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.07, 1.52], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 46% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 93% reduced risk up to 52% increased risk.

Figure C13D. Adult (18+) 1-month patient death HR estimate (deceased donor grafts)





Figure C14D. Adult (18+) 1-month patient death HR program comparison (deceased donor grafts)





C. Transplant Information

Table C11L. Adult (18+) 1-month patient survival (living donor graft recipients)
Single organ transplants performed between 07/01/2017 and 12/31/2019
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	112	14,136
Estimated probability of surviving at 1 month (unadjusted for patient and donor characteristics)	100.00%	99.78%
Expected probability of surviving at 1 month (adjusted for patient and donor characteristics)	99.77%	--
Number of observed deaths during the first month after transplant	0	31
Number of expected deaths during the first month after transplant	0.26	--
Estimated hazard ratio*	0.89	--
95% credible interval for the hazard ratio**	[0.11, 2.47]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.11, 2.47], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 11% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 89% reduced risk up to 147% increased risk.

Figure C13L. Adult (18+) 1-month patient death HR estimate (living donor grafts)





Figure C14L. Adult (18+) 1-month patient death HR program comparison (living donor grafts)





C. Transplant Information

Table C12. Adult (18+) 1-year patient survival
Single organ transplants performed between 07/01/2017 and 12/31/2019
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	389	44,658
Estimated probability of surviving at 1 year (unadjusted for patient and donor characteristics)	98.67%	97.61%
Expected probability of surviving at 1 year (adjusted for patient and donor characteristics)	97.31%	--
Number of observed deaths during the first year after transplant	5	943
Number of expected deaths during the first year after transplant	9.23	--
Estimated hazard ratio*	0.62	--
95% credible interval for the hazard ratio**	[0.25, 1.16]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.25, 1.16], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 38% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 75% reduced risk up to 16% increased risk.

Figure C15. Adult (18+) 1-year patient death HR estimate





Figure C16. Adult (18+) 1-year patient death HR program comparison





C. Transplant Information

Table C12D. Adult (18+) 1-year patient survival (deceased donor graft recipients)
Single organ transplants performed between 07/01/2017 and 12/31/2019
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	277	30,522
Estimated probability of surviving at 1 year (unadjusted for patient and donor characteristics)	98.88%	96.96%
Expected probability of surviving at 1 year (adjusted for patient and donor characteristics)	96.66%	--
Number of observed deaths during the first year after transplant	3	818
Number of expected deaths during the first year after transplant	8.23	--
Estimated hazard ratio*	0.49	--
95% credible interval for the hazard ratio**	[0.16, 1.00]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.16, 1.00], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 51% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 84% reduced risk up to 0% increased risk.

Figure C15D. Adult (18+) 1-year patient death HR estimate (deceased donor grafts)





Figure C16D. Adult (18+) 1-year patient death HR program comparison (deceased donor grafts)





C. Transplant Information

Table C12L. Adult (18+) 1-year patient survival (living donor graft recipients)
Single organ transplants performed between 07/01/2017 and 12/31/2019
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	112	14,136
Estimated probability of surviving at 1 year (unadjusted for patient and donor characteristics)	98.14%	99.01%
Expected probability of surviving at 1 year (adjusted for patient and donor characteristics)	98.95%	--
Number of observed deaths during the first year after transplant	2	125
Number of expected deaths during the first year after transplant	0.99	--
Estimated hazard ratio*	1.34	--
95% credible interval for the hazard ratio**	[0.36, 2.93]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.36, 2.93], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 34% higher risk of patient death compared to an average program, but NYUC's performance could plausibly range from 64% reduced risk up to 193% increased risk.

Figure C15L. Adult (18+) 1-year patient death HR estimate (living donor grafts)





Figure C16L. Adult (18+) 1-year patient death HR program comparison (living donor grafts)





C. Transplant Information

Table C13. Adult (18+) 3-year patient survival
Single organ transplants performed between 01/01/2015 and 06/30/2017
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	185	36,746
Estimated probability of surviving at 3 years (unadjusted for patient and donor characteristics)	93.51%	94.00%
Expected probability of surviving at 3 years (adjusted for patient and donor characteristics)	91.18%	--
Number of observed deaths during the first 3 years after transplant	12	2,189
Number of expected deaths during the first 3 years after transplant	16.26	--
Estimated hazard ratio*	0.77	--
95% credible interval for the hazard ratio**	[0.42, 1.22]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.42, 1.22], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 23% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 58% reduced risk up to 22% increased risk.

Figure C17. Adult (18+) 3-year patient death HR estimate





Figure C18. Adult (18+) 3-year patient death HR program comparison





C. Transplant Information

Table C13D. Adult (18+) 3-year patient survival (deceased donor graft recipients)
Single organ transplants performed between 01/01/2015 and 06/30/2017
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	157	24,671
Estimated probability of surviving at 3 years (unadjusted for patient and donor characteristics)	92.36%	92.54%
Expected probability of surviving at 3 years (adjusted for patient and donor characteristics)	89.99%	--
Number of observed deaths during the first 3 years after transplant	12	1,829
Number of expected deaths during the first 3 years after transplant	15.65	--
Estimated hazard ratio*	0.79	--
95% credible interval for the hazard ratio**	[0.43, 1.26]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.43, 1.26], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 21% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 57% reduced risk up to 26% increased risk.

Figure C17D. Adult (18+) 3-year patient death HR estimate (deceased donor grafts)





Figure C18D. Adult (18+) 3-year patient death HR program comparison (deceased donor grafts)





C. Transplant Information

Table C13L. Adult (18+) 3-year patient survival (living donor graft recipients)
Single organ transplants performed between 01/01/2015 and 06/30/2017
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	28	12,075
Estimated probability of surviving at 3 years (unadjusted for patient and donor characteristics)	100.00%	96.99%
Expected probability of surviving at 3 years (adjusted for patient and donor characteristics)	97.85%	--
Number of observed deaths during the first 3 years after transplant	0	360
Number of expected deaths during the first 3 years after transplant	0.61	--
Estimated hazard ratio*	0.77	--
95% credible interval for the hazard ratio**	[0.09, 2.14]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.09, 2.14], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 23% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 91% reduced risk up to 114% increased risk.

Figure C17L. Adult (18+) 3-year patient death HR estimate (living donor grafts)





Figure C18L. Adult (18+) 3-year patient death HR program comparison (living donor grafts)





C. Transplant Information

Table C14. Pediatric (<18) 1-month patient survival
Single organ transplants performed between 07/01/2017 and 12/31/2019
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	6	1,938
Estimated probability of surviving at 1 month (unadjusted for patient and donor characteristics)	100.00%	99.79%
Expected probability of surviving at 1 month (adjusted for patient and donor characteristics)	99.77%	--
Number of observed deaths during the first month after transplant	0	4
Number of expected deaths during the first month after transplant	0.01	--
Estimated hazard ratio*	0.99	--
95% credible interval for the hazard ratio**	[0.12, 2.77]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.12, 2.77], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 1% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 88% reduced risk up to 177% increased risk.

Figure C19. Pediatric (<18) 1-month patient death HR estimate





Figure C20. Pediatric (<18) 1-month patient death HR program comparison





C. Transplant Information

Table C14D. Pediatric (<18) 1-month patient survival (deceased donor graft recipients)
Single organ transplants performed between 07/01/2017 and 12/31/2019
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	3	1,290
Estimated probability of surviving at 1 month (unadjusted for patient and donor characteristics)	100.00%	99.84%
Expected probability of surviving at 1 month (adjusted for patient and donor characteristics)	99.85%	--
Number of observed deaths during the first month after transplant	0	2
Number of expected deaths during the first month after transplant	0.00	--
Estimated hazard ratio*	1.00	--
95% credible interval for the hazard ratio**	[0.12, 2.78]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.12, 2.78], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 0% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 88% reduced risk up to 178% increased risk.

Figure C19D. Pediatric (<18) 1-month patient death HR estimate (deceased donor grafts)





Figure C20D. Pediatric (<18) 1-month patient death HR program comparison (deceased donor grafts)





C. Transplant Information

Table C14L. Pediatric (<18) 1-month patient survival (living donor graft recipients)
Single organ transplants performed between 07/01/2017 and 12/31/2019
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	3	648
Estimated probability of surviving at 1 month (unadjusted for patient and donor characteristics)	100.00%	99.69%
Expected probability of surviving at 1 month (adjusted for patient and donor characteristics)	99.69%	--
Number of observed deaths during the first month after transplant	0	2
Number of expected deaths during the first month after transplant	0.01	--
Estimated hazard ratio*	1.00	--
95% credible interval for the hazard ratio**	[0.12, 2.77]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.12, 2.77], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 0% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 88% reduced risk up to 177% increased risk.

Figure C19L. Pediatric (<18) 1-month patient death HR estimate (living donor grafts)





Figure C20L. Pediatric (<18) 1-month patient death HR program comparison (living donor grafts)





C. Transplant Information

Table C15. Pediatric (<18) 1-year patient survival
Single organ transplants performed between 07/01/2017 and 12/31/2019
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	6	1,938
Estimated probability of surviving at 1 year (unadjusted for patient and donor characteristics)	100.00%	99.61%
Expected probability of surviving at 1 year (adjusted for patient and donor characteristics)	99.63%	--
Number of observed deaths during the first year after transplant	0	7
Number of expected deaths during the first year after transplant	0.02	--
Estimated hazard ratio*	0.99	--
95% credible interval for the hazard ratio**	[0.12, 2.76]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.12, 2.76], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 1% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 88% reduced risk up to 176% increased risk.

Figure C21. Pediatric (<18) 1-year patient death HR estimate





Figure C22. Pediatric (<18) 1-year patient death HR program comparison





C. Transplant Information

Table C15D. Pediatric (<18) 1-year patient survival (deceased donor graft recipients)
Single organ transplants performed between 07/01/2017 and 12/31/2019
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	3	1,290
Estimated probability of surviving at 1 year (unadjusted for patient and donor characteristics)	100.00%	99.56%
Expected probability of surviving at 1 year (adjusted for patient and donor characteristics)	99.57%	--
Number of observed deaths during the first year after transplant	0	5
Number of expected deaths during the first year after transplant	0.01	--
Estimated hazard ratio*	0.99	--
95% credible interval for the hazard ratio**	[0.12, 2.77]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.12, 2.77], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 1% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 88% reduced risk up to 177% increased risk.

Figure C21D. Pediatric (<18) 1-year patient death HR estimate (deceased donor grafts)





Figure C22D. Pediatric (<18) 1-year patient death HR program comparison (deceased donor grafts)





C. Transplant Information

Table C15L. Pediatric (<18) 1-year patient survival (living donor graft recipients)
Single organ transplants performed between 07/01/2017 and 12/31/2019
Retransplants excluded

	NYUC	U.S.
Number of transplants evaluated	3	648
Estimated probability of surviving at 1 year (unadjusted for patient and donor characteristics)	100.00%	99.69%
Expected probability of surviving at 1 year (adjusted for patient and donor characteristics)	99.69%	--
Number of observed deaths during the first year after transplant	0	2
Number of expected deaths during the first year after transplant	0.01	--
Estimated hazard ratio*	1.00	--
95% credible interval for the hazard ratio**	[0.12, 2.77]	--

* The hazard ratio provides an estimate of how New York University Medical Center (NYUC)'s results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYUC's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.12, 2.77], indicates the location of NYUC's true hazard ratio with 95% probability. The best estimate is 0% lower risk of patient death compared to an average program, but NYUC's performance could plausibly range from 88% reduced risk up to 177% increased risk.

Figure C21L. Pediatric (<18) 1-year patient death HR estimate (living donor grafts)





Figure C22L. Pediatric (<18) 1-year patient death HR program comparison (living donor grafts)





C. Transplant Information

Table C16. Pediatric (<18) 3-year patient survival

Single organ transplants performed between 01/01/2015 and 06/30/2017
Retransplants excluded

This center did not perform any
transplants relevant to
this table during
01/01/2015-06/30/2017

Figure C23. Pediatric (<18) 3-year patient death HR estimate

This center did not perform any
transplants relevant to
this figure during
01/01/2015-06/30/2017

Figure C24. Pediatric (<18) 3-year patient death HR program comparison

This center did not perform any
transplants relevant to
this figure during
01/01/2015-06/30/2017



C. Transplant Information

Table C16D. Pediatric (<18) 3-year patient survival (deceased donor graft recipients)
Single organ transplants performed between 01/01/2015 and 06/30/2017
Retransplants excluded

This center did not perform any
transplants relevant to
this table during
01/01/2015-06/30/2017

**Figure C23D. Pediatric (<18)
3-year patient death HR estimate
(deceased donor grafts)**

This center did not perform any
transplants relevant to
this figure during
01/01/2015-06/30/2017

**Figure C24D. Pediatric (<18) 3-year patient death HR program
comparison (deceased donor grafts)**

This center did not perform any
transplants relevant to
this figure during
01/01/2015-06/30/2017



C. Transplant Information

Table C16L. Pediatric (<18) 3-year patient survival (living donor graft recipients)
Single organ transplants performed between 01/01/2015 and 06/30/2017
Retransplants excluded

This center did not perform any
transplants relevant to
this table during
01/01/2015-06/30/2017

**Figure C23L. Pediatric (<18)
3-year patient death HR estimate
(living donor grafts)**

This center did not perform any
transplants relevant to
this figure during
01/01/2015-06/30/2017

**Figure C24L. Pediatric (<18) 3-year patient death HR program
comparison (living donor grafts)**

This center did not perform any
transplants relevant to
this figure during
01/01/2015-06/30/2017



C. Transplant Information

Table C17. Multi-organ transplant graft survival: 07/01/2017 - 12/31/2019

Adult (18+) Transplants

Transplant Type	First-Year Outcomes					
	Transplants Performed		Kidney Graft Failures		Estimated Kidney Graft Survival	
	NYUC-TX1	USA	NYUC-TX1	USA	NYUC-TX1	USA
Kidney-Heart	8	507	1	76	87.5%	84.7%
Kidney-Liver	18	1,691	1	179	94.4%	88.9%
Kidney-Pancreas	12	2,109	0	76	100.0%	96.2%

Pediatric (<18) Transplants

No pediatric (<18) multi-organ transplants were performed

Table C18. Multi-organ transplant patient survival: 07/01/2017 - 12/31/2019

Adult (18+) Transplants

Transplant Type	First-Year Outcomes					
	Transplants Performed		Patient Deaths		Estimated Patient Survival	
	NYUC-TX1	USA	NYUC-TX1	USA	NYUC-TX1	USA
Kidney-Heart	8	507	0	58	100.0%	88.3%
Kidney-Liver	18	1,691	1	147	94.4%	90.8%
Kidney-Pancreas	12	2,109	0	46	100.0%	97.6%

Pediatric (<18) Transplants

No pediatric (<18) multi-organ transplants were performed



D. Living Donor Information

Table D1. Living donor summary: 07/01/2017 - 06/30/2020

Living Donor Follow-Up	This Center			United States		
	07/2017-06/2018	07/2018-06/2019	07/2019-12/2019	07/2017-06/2018	07/2018-06/2019	07/2019-12/2019
Number of Living Donors	38	57	33	6,151	6,685	3,483
6-Month Follow-Up						
Donors due for follow-up	38	57	33	6,150	6,682	3,468
Timely clinical data	27 71.1%	44 77.2%	29 87.9%	5,395 87.7%	5,832 87.3%	2,529 72.9%
Timely lab data	21 55.3%	39 68.4%	16 48.5%	5,213 84.8%	5,629 84.2%	2,252 64.9%
12-Month Follow-Up						
Donors due for follow-up	38	57		6,148	6,681	
Timely clinical data	22 57.9%	43 75.4%		5,095 82.9%	5,075 76.0%	
Timely lab data	17 44.7%	27 47.4%		4,818 78.4%	4,562 68.3%	
24-Month Follow-Up						
Donors due for follow-up	38			6,147		
Timely clinical data	27 71.1%			4,134 67.3%		
Timely lab data	10 26.3%			3,575 58.2%		

Follow-up completion standards through 2 years post-donation were implemented in policy on February 1, 2013.