



COVID-19 Guide

Adjustments to Transplant Program and OPO Evaluation Metrics

The Scientific Registry of Transplant Recipients (SRTR), under contract from the Health Resources and Services Administration (HRSA), is charged with evaluating the performance of the nation's transplant system through publication of semi-annual transplant program-specific reports (PSRs) and organ procurement organization (OPO)-specific reports (OSRs). These reports contain performance metrics covering various time periods. For OPOs, these metrics include eligible death conversion rates and deceased donor organ yield. For transplant programs, they include pre-transplant mortality rates (formerly called waitlist mortality rates), transplant rates, organ offer acceptance rates, patient mortality after listing, and 1-month, 90-day, 1-year, 1-year conditional on 90-day, and 3-year posttransplant outcomes including graft survival and patient survival.

In response to the current global pandemic, SRTR modified the evaluation metrics for transplant programs and OPOs for the reports released in January 2021, July 2021, January 2022 and July 2022. These reports made adjustments to transplant program and OPO performance metrics so that data beyond the declaration of a national public health emergency on March 13, 2020, were not included in the metrics.

Modifications for the January 2023 reporting cycle were considered at the Analytic Methods Subcommittee of the SRTR Review Committee (SRC) at its meeting on March 24, 2021, and the full SRC meetings April 27, 2021 and on January 11, 2022. Both the Analytic Methods Subcommittee and the full SRC recommended an ongoing carve out of the first quarter of the pandemic (March 13, 2020 through June 12, 2020) from adjusted performance metrics, as detailed below. These recommendations were reviewed by HRSA's Division of Transplantation, which oversees SRTR. HRSA approved these recommendations, which SRTR will implement for the January 2023 reporting cycle. These changes will remain in force beyond the January 2023 reporting cycle, unless otherwise amended:

Posttransplant Outcomes (including 1-month, 90-day, 1-year, 1-year conditional on 90-day, and 3-year graft and patient survival): Evaluation cohorts will exclude transplants performed between March 13, 2020 and June 12, 2020, inclusive of March 13 and June 12. Patients given transplants before March 13, 2020 will have follow-up censored on March 12, 2020. Patients given transplants after June 12, 2020 will resume normal follow-up. Follow-up will not resume for patients given transplants before March 13, 2020 who are alive with function on June 12, 2020; however, this may be reconsidered as SRTR continues to explore moving to a period-prevalent methodology:

1-month, 90-day, 1-year & 1-year conditional on 90-day Patient and Graft Survival Evaluations: Transplants 7/1/2019-3/12/2020, follow-up through 3/12/2020. Transplants 6/13/2020-12/31/2021, follow-up through 6/30/2022.

3-year Patient and Graft Survival Evaluations: Transplants 1/1/2017-6/30/2019; follow-up through 3/12/2020.

Pre-Transplant Mortality Rate (formerly called Waitlist Mortality Rate): These evaluations are based on normal reporting cohorts.



COVID-19 Guide

Days after listing (and before transplant) between 7/1/2020 and 6/30/2022.

Transplant Rate: These evaluations are based on normal reporting cohorts.

Candidates on the waitlist 7/1/2020-6/30/2022.

Overall Rate of Mortality After Listing: These evaluations are based on normal reporting cohorts.

Evaluation period: 7/1/2020-6/30/2022.

Offer Acceptance Rate: These evaluations are based on normal reporting cohorts.

Offers received 1/1/2021-12/31/2021.

These decisions will apply to the evaluations released in the SRTR's semi-annual program-specific reports scheduled for release on January 6, 2023. These changes have been communicated to the leadership of the Organ Procurement and Transplantation Network's (OPTN) Membership and Professional Standards Committee (MSPC). These decisions will then be re-evaluated as more information becomes available in preparation for the release scheduled for July 2023.

As with the July 2022 reports, SRTR will continue to report descriptive data beyond March 12, 2020, e.g., waitlist counts, transplant counts, recipient characteristics, donor counts, donor characteristics, etc., but will alter data for performance evaluation metrics as described above.



User Guide

This report contains a wide range of useful information about the heart transplant program at NY Presbyterian Hospital/Columbia Univ. Medical Center. The report has three main sections:

- A. Program Summary
- B. Waiting List Information
- C. Transplant Information

The Program Summary is a one-page summary highlighting characteristics of the program, including the number of candidates on the waiting list, the number of transplants performed at the program, the number of patients being cared for by the program, and patient outcomes, including outcomes while on the waiting list (the transplant rate and the death rate while on the waiting list) and outcomes after transplant (patient and graft survival probabilities). If the program performed transplants in both adults and children, survival probabilities for adults and children (pediatrics) are provided separately. For each of the outcomes measures presented, a comparison is provided showing what would be expected at this program if it were performing as similar programs around the country perform when treating similar patients. More details regarding these outcome measures are provided in Sections B and C of the report.

The Waiting List Information section contains more detailed information on how many candidates are on the waiting list at the program, the types of candidates on the waiting list, how long candidates typically have to wait for a transplant at this program, how frequently candidates successfully receive a transplant, and how often candidates on the waiting list die before receiving a transplant.

Table B1 shows the activity on this program's waiting list during two recent 1-year periods and provides comparisons to all programs within this program's OPTN region (see <http://optn.transplant.hrsa.gov/members/regions.asp> for information on OPTN regions) and the nation as a whole. Tables B2 and B3 describe the candidates on the waiting list at this program, with comparisons to candidates waiting in the same donor service area (OPO/DSA) the OPTN region, and the nation as a whole.

Table B4 shows how many candidates were removed from the waiting list because they received a transplant. The program's transplant rate is calculated as the number of candidates who received a transplant divided by the person-years observed at the program (person-years is a combination of how many candidates were on the waiting list along with how long each candidate was followed since some candidates are not on the waiting list for the entire year). The transplant rate and comparisons to what would be expected at this program are presented in Figures B1 and B2. Figure B1 shows the transplant rate compared to what was expected at this program. The expected transplant rate is an estimate of what we would expect at this program if it were performing transplants at rates similar to other programs in the US with similar candidates on their waiting lists. The expected rate is only an estimate, and is made with a certain level of uncertainty. This uncertainty is shown in Figure B2. Figure B2 displays the ratio of the observed to the expected transplant rate. A ratio of 1 indicates that the observed transplant rate was equal to the expected transplant rate, while a ratio less than 1 indicates the observed rate was lower than expected rate and a ratio greater than 1 indicates the observed rate was higher than the expected rate. However, the level of uncertainty must be considered when interpreting these numbers. The 95% interval is also shown on Figure B2. This interval provides a range within which the true ratio of observed to expected transplant rates is likely to be. If this



User Guide

confidence interval includes (crosses) 1.0, then we cannot say that this program's observed transplant rate is different from what would be expected. The observed deceased donor transplant rate at this program was 85.4 per 100 person-years. Transplant rates are also provided for adult and pediatric patients separately along with comparisons to adult and pediatric rates in the DSA, the OPTN region, and the nation. Please refer to the PSR Technical Methods documentation available at <http://www.srtr.org> for more detail regarding how expected rates are calculated.

The pre-transplant mortality rate (previously called the waiting list mortality rate) for candidates on the waiting list is presented in Table B5 and Figures B4-B6. These data are presented in the same way as the transplant rate data in the previous section. The intent of this table and figures is to describe risk of death once candidates are listed rather than while they are listed, but before they are transplanted. Therefore, time at risk and deaths after removal from the waiting list for reasons other than transplant, transfer to another transplant program, or recovery (no longer needing a transplant), and before any subsequent transplant, are included. As with transplant rates, mortality rates should be interpreted carefully taking into consideration the interval displayed in Figure B5. For a complete description of how observed and expected mortality rates are calculated, please refer to the technical documentation available at <http://www.srtr.org>.

Survival from listing is presented in Table B6 and Figures B7-B9. These data are presented in the same way as the pre-transplant mortality rate data in the previous section. The intent of this table and figures is to describe risk of death once candidates are listed rather than while they are listed, including after a transplant. As with transplant rates, mortality rates should be interpreted carefully taking into consideration the interval displayed in Figure B8. For a complete description of how observed and expected mortality rates are calculated, please refer to the technical documentation available at <http://www.srtr.org>.

Table B7 presents information on what happens to candidates on the waiting list by three different time points after listing: 6 months, 12 months, and 18 months. The table displays percentages of candidates who have died, been removed from the waiting list, been transplanted, or been transferred or lost-to-follow-up. Tables B8 and B9 provide more detail regarding how many candidates have received a deceased donor transplant by certain time points during the first 3 years after being put on the transplant waiting list. Each row of Tables B8 and B9 presents the percent of candidates who received a deceased donor transplant by each time point. Table B10 presents data on the time it took for different percentages of patients to be transplanted for candidates added to the list between 07/01/2016 and 12/31/2021. The time it took for 5% (the 5th percentile) of patients to receive a transplant at this program was 0.2 months. If "Not Observed" is displayed in the table, then too few candidates received transplants before 06/30/2022 to calculate a particular percentile of transplant times.

Table B11 contains a summary of the offer acceptance practices of the program. The offer acceptance ratio indicates whether the program is more or less likely to accept offers than the average program. If the offer acceptance ratio is greater than 1.0, then the program tends to accept more offers than average; if the offer acceptance ratio is less than 1.0, then the program tends to accept fewer offers than average. Figure B10 shows the distribution of program offer acceptance rates as well as the offer acceptance rate for this program. Figures B11 - B15 similarly show offer acceptance rates for subsets of offers.



User Guide

The Transplant Information section begins with descriptions of transplant recipients in Tables C1 and C2. Data on recipients of deceased donor transplants are presented (Tables C1D and C2D); if applicable, data on recipients of living donor transplants are presented separately (Tables C1L and C2L). Comparisons to the region and the nation as a whole are provided. A description of the deceased donors used at this program is provided in Table C3D, along with characteristics of living donors in Table C3L, if applicable. Finally, information on the transplant procedure for deceased and living donor transplants is presented in Tables C4D and C4L, respectively.

Starting with Table C5, transplant outcomes are presented along with comparisons to what would be expected at this program and what happened in the nation as a whole. Tables C5-C14 (tables C5-C10 for Pancreas) present information on graft survival (survival of the transplanted organ), with data presented separately for adult and pediatric recipients. Patients are followed from the time of transplant until either failure of the transplanted organ or death, whichever comes first. Please refer to the technical methods for more information on these calculations (<http://www.srtr.org>).

While Tables C5-C14 present data on graft survival, Tables C15-C20 (tables C11-C20 for Pancreas) present information on patient survival. For these tables, patients are followed from the time of transplant until death, regardless of whether the transplant is functioning or the patient required another transplant to survive.

Tables C21 and C22 summarize the multiorgan transplant outcomes at this program. The summary statistics in these tables are descriptive and are not risk-adjusted for different donor and candidate characteristics.

Additional information regarding the technical methods and the risk adjustment models used to estimate expected event rates is available on the SRTR website at <http://www.srtr.org>. We welcome and encourage feedback on these reports. Please feel free to share feedback with the SRTR at the following e-mail: srtr@srtr.org.



Table of Contents

Section	Page
COVID-19 Guide	i
User Guide	iii
A. Program Summary	
Program Summary	1
B. Waiting List Information	
Waiting list activity	2
Demographic characteristics of waiting list candidates	3
Medical characteristics of waiting list candidates	4
Deceased donor transplant rates	5
Pre-transplant mortality rates (formerly called Waiting list mortality rates)	6
Patient survival from listing	7
Waiting list candidate status after listing	8
Percent of candidates with deceased donor transplants: demographic characteristics	9
Percent of candidates with deceased donor transplants: medical characteristics	10
Time to transplant for waiting list candidates	11
Offer acceptance practices	12
C. Transplant Information	
Deceased donor transplant recipient demographic characteristics	14
Deceased donor transplant recipient medical characteristics	15
Deceased donor characteristics	16
Deceased donor transplant characteristics	17
Deceased donor graft survival	18
Deceased donor patient survival	28
Multi-organ transplant graft survival	34
Multi-organ transplant patient survival	34



A. Program Summary

Figure A1. Waiting list and transplant activity




Table A1. Census of transplant recipients

Recipients	07/01/2020-06/30/2021	07/01/2021-06/30/2022
Transplanted at this center	95	106
Followed by this center*	876	899
...transplanted at this program	831	854
...transplanted elsewhere	45	45

* Recipients followed are transplant recipients for whom the center has submitted a post-transplant follow-up form for a transplant that took place before the 12-month interval for each column.

Figure A2. Transplant rates
07/01/2020 - 06/30/2022




Figure A3. Pre-transplant mortality rates
07/01/2020 - 06/30/2022




Figure A4. First-year adult graft and patient survival:
07/01/2019 - 03/12/2020, 06/13/2020 - 12/31/2021





Figure A5. First-year pediatric graft & patient survival:
07/01/2019 - 03/12/2020, 06/13/2020 - 12/31/2021





B. Waiting List Information

Table B1. Waiting list activity summary: 07/01/2020 - 06/30/2022

Waiting List Registrations	Counts for this center		Activity for 07/01/2021 to 06/30/2022 as percent of registrants on waiting list on 07/01/2021		
	07/01/2020-06/30/2021	07/01/2021-06/30/2022	This Center (%)	OPTN Region (%)	U.S. (%)
On waiting list at start	103	127	100.0	100.0	100.0
Additions					
New listings at this center	143	133	104.7	116.1	133.8
Removals					
Transferred to another center	4	2	1.6	2.1	2.2
Received living donor transplant*	0	0	0.0	0.0	0.0
Received deceased donor transplant*	95	106	83.5	95.3	105.0
Died	7	7	5.5	5.9	6.5
Transplanted at another center	0	1	0.8	1.5	0.9
Deteriorated	6	6	4.7	4.7	6.6
Recovered	2	7	5.5	6.2	6.7
Other reasons	5	7	5.5	11.1	10.5
On waiting list at end of period	127	124	97.6	89.4	95.4

* These patients were removed from waiting list with removal code indicating transplant; this may not equal the number of transplants performed at this center during the specified period.



B. Waiting List Information

Table B2. Demographic characteristics of waiting list candidates**Candidates registered on the waiting list between 07/01/2021 and 06/30/2022**

Demographic Characteristic	New Waiting List Registrations 07/01/2021 to 06/30/2022 (%)			All Waiting List Registrations on 06/30/2022 (%)		
	This Center (N=133)	OPTN Region (N=396)	U.S. (N=4,858)	This Center (N=124)	OPTN Region (N=305)	U.S. (N=3,465)
All (%)	100.0	100.0	100.0	100.0	100.0	100.0
Ethnicity/Race (%)*						
White	39.8	47.0	56.6	41.9	46.9	55.8
African-American	23.3	26.5	25.1	28.2	26.9	28.2
Hispanic/Latino	22.6	19.4	12.8	21.0	20.0	11.7
Asian	14.3	6.8	4.4	8.9	6.2	3.1
Other	0.0	0.3	1.2	0.0	0.0	1.2
Unknown	0.0	0.0	0.0	0.0	0.0	0.0
Age (%)						
<2 years	9.8	4.3	5.1	6.5	3.0	4.1
2-11 years	7.5	3.8	4.6	1.6	1.0	5.9
12-17 years	6.0	3.3	4.6	1.6	1.6	4.0
18-34 years	8.3	8.6	9.7	8.1	7.5	10.2
35-49 years	15.0	17.2	17.3	24.2	24.6	22.0
50-64 years	36.8	44.4	41.2	42.7	46.2	42.2
65-69 years	13.5	13.6	14.4	12.1	14.1	10.7
70+ years	3.0	4.8	3.1	3.2	2.0	0.9
Gender (%)						
Male	66.9	72.2	70.8	71.0	73.8	74.8
Female	33.1	27.8	29.2	29.0	26.2	25.2

* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



B. Waiting List Information

Table B3. Medical characteristics of waiting list candidates

Candidates registered on the waiting list between 07/01/2021 and 06/30/2022

Medical Characteristic	New Waiting List Registrations 07/01/2021 to 06/30/2022 (%)			All Waiting List Registrations on 06/30/2022 (%)		
	This Center (N=133)	OPTN Region (N=396)	U.S. (N=4,858)	This Center (N=124)	OPTN Region (N=305)	U.S. (N=3,465)
All (%)	100.0	100.0	100.0	100.0	100.0	100.0
Blood Type (%)						
O	51.1	47.0	44.7	58.1	56.7	60.7
A	27.1	30.8	36.7	26.6	26.2	27.2
B	18.8	18.2	14.7	14.5	15.1	10.5
AB	3.0	4.0	4.0	0.8	2.0	1.5
Unknown	0.0	0.0	0.0	0.0	0.0	0.0
Previous Transplant (%)						
Yes	8.3	5.1	4.0	7.3	4.6	4.0
No	91.7	94.9	96.0	92.7	95.4	96.0
Unknown	0.0	0.0	0.0	0.0	0.0	0.0
Primary Disease (%)						
Cardiomyopathy	55.6	59.1	58.4	55.6	56.4	56.2
Coronary Artery Disease	16.5	23.0	23.1	25.8	28.2	24.2
Retransplant/Graft Failure	4.5	2.3	3.4	4.0	2.6	3.2
Valvular Heart Disease	1.5	1.3	1.0	4.0	2.0	0.8
Congenital Heart Disease	20.3	9.3	11.5	9.7	6.2	13.4
Other	1.5	5.1	2.6	0.8	4.6	2.1
Missing	0.0	0.0	0.0	0.0	0.0	0.0
Medical Urgency Status at Listing (%)						
Status 1A	21.1	9.3	8.6	8.9	5.2	5.3
Status 1B	1.5	1.0	2.8	4.0	5.2	7.9
Status 2	0.8	1.0	2.7	11.3	8.5	11.8
Adult Status 1	6.8	5.6	4.7	0.0	0.7	0.5
Adult Status 2	10.5	31.6	24.4	2.4	5.6	4.6
Adult Status 3	1.5	4.5	8.6	0.8	3.3	4.1
Adult Status 4	33.1	28.5	29.9	38.7	39.7	38.5
Adult Status 5	4.5	4.3	2.7	1.6	2.6	3.6
Adult Status 6	20.3	13.6	14.1	32.3	28.5	21.3
Temporarily Inactive	0.0	0.5	1.4	0.0	0.7	2.4



B. Waiting List Information

Table B4D. Deceased donor transplant rates: 07/01/2020 - 06/30/2022

Waiting List Registrations	This Center	OPO/DSA	Region	U.S.
All Candidates				
Count on waiting list at start*	103	292	323	3,514
Person Years**	235.5	585.0	641.2	7,056.5
Removals for Transplant	201	585	617	7,652
Adult (18+) Candidates				
Count on waiting list at start*	91	274	305	3,079
Person Years**	213.9	554.2	610.4	6,159.0
Removals for transplant	152	525	557	6,651
Pediatric (<18) Candidates				
Count on waiting list at start*	12	18	18	435
Person Years**	21.6	30.9	30.9	897.5
Removals for transplant	49	60	60	1,001

* Counts in this table may be lower than similar counts in other waiting list tables, such as Table B1. A small percentage (~1%) of patients are found to have died or been transplanted before being removed from the waiting list, so these patients are excluded if the event occurs prior to the start of the study period. Inactive time on the waiting list is included in the calculations for this table.

** Person years are calculated as days (converted to fractional years). The number of days from July 1 or from the date of first wait listing until death, transplant, removal from the waiting list or June 30.

Figure B1D. Observed and expected deceased donor transplant rates: 07/01/2020 - 06/30/2022




Figure B2D. Deceased donor transplant rate ratio estimate





Figure B3D. Observed adult (18+) and pediatric (<18) deceased donor transplant rates: 07/01/2020 - 06/30/2022





B. Waiting List Information

Table B5. Pre-transplant mortality rates: 07/01/2020 - 06/30/2022

Waiting List Registrations	This Center	OPO/DSA	Region	U.S.
All Candidates				
Count on waiting list at start*	103	292	323	3,514
Person Years**	253.8	644.3	719.4	8,037.2
Number of deaths	16	47	54	693
Adult (18+) Candidates				
Count on waiting list at start*	91	274	305	3,079
Person Years**	230.3	605.0	680.1	7,033.8
Number of deaths	8	37	44	584
Pediatric (<18) Candidates				
Count on waiting list at start*	12	18	18	435
Person Years**	23.5	39.3	39.3	1,003.4
Number of deaths	8	10	10	109

* Counts in this table may be lower than similar counts in other waiting list tables, such as Table B1. A small percentage (~1%) of patients are found to have died or been transplanted before being removed from the waiting list, so these patients are excluded if the event occurs prior to the start of the study period. Inactive time on the waiting list is included in the calculations for this table.

** Person years are calculated as days (converted to fractional years). The number of days from July 1 or from the date of first wait listing until death, transplant, 60 days after recovery, transfer or June 30.

Figure B4. Observed and expected pre-transplant mortality rates: 07/01/2020 - 06/30/2022




Figure B6. Observed adult (18+) and pediatric (<18) pre-transplant mortality rates: 07/01/2020 - 06/30/2022





Figure B5. Pre-transplant mortality rate ratio estimate





B. Waiting List Information

Table B6. Rates of patient mortality after listing: 07/01/2020 - 06/30/2022

Waiting List Registrations	This Center	OPO/DSA	Region	U.S.
All Patients				
Count at risk during the evaluation period	704	1,886	2,056	28,175
Person-years*	950.5	2,540.5	2,777.1	38,163.1
Number of Deaths	37	138	152	2,306
Adult (18+) Patients				
Count at risk during the evaluation period	519	1,644	1,814	24,127
Person-years*	697.7	2,217.8	2,454.4	32,646.6
Number of Deaths	22	118	132	2,055
Pediatric (<18) Patients				
Count at risk during the evaluation period	185	242	242	4,048
Person-years*	252.8	322.7	322.7	5,516.5
Number of Deaths	15	20	20	251

* Person-years are calculated as days (converted to fractional years). The number of days from 07/01/2020, or from the date of first wait listing until death, reaching 5 years after listing or June 30, 2022.

** Patient mortality after listing describes the relative survival experience of patients after listing. It depends on many factors, some of which are outside of the control of the transplant program. For example, availability of organs may not be the same in every part of the country.

Figure B7. Observed and expected rates of patient mortality after listing: 07/01/2020 - 06/30/2022




Figure B8. HR estimate of patient mortality after listing





Figure B9. Observed adult (18+) and pediatric (<18) rates of patient mortality after listing: 07/01/2020 - 06/30/2022





B. Waiting List Information

Table B7. Waiting list candidate status after listing

Candidates registered on waiting list between 01/01/2020 and 12/31/2020

Waiting list status (survival status)	This Center (N=114)			U.S. (N=4,594)		
	Months Since Listing			Months Since Listing		
	6	12	18	6	12	18
Alive on waiting list (%)	32.5	21.9	13.2	30.2	19.8	14.3
Died on the waiting list without transplant (%)	3.5	3.5	4.4	3.1	3.7	4.0
Removed without transplant (%):						
Condition worsened (status unknown)	1.8	1.8	1.8	3.2	4.0	4.4
Condition improved (status unknown)	0.0	0.9	0.9	0.6	1.5	2.4
Refused transplant (status unknown)	0.0	0.0	0.0	0.1	0.2	0.2
Other	1.8	2.6	4.4	1.8	2.6	3.5
Transplant (living or deceased donor) (%):						
Functioning (alive)	57.9	60.5	36.8	56.7	58.4	39.1
Failed-Retransplanted (alive)	0.0	0.0	0.0	0.2	0.2	0.3
Failed-alive not retransplanted	0.0	0.0	0.0	0.0	0.0	0.0
Died	2.6	6.1	7.0	3.3	5.2	6.7
Status Yet Unknown*	0.0	2.6	31.6	0.3	3.7	24.2
Lost or Transferred (status unknown) (%)	0.0	0.0	0.0	0.4	0.7	0.8
TOTAL (%)	100.0	100.0	100.0	100.0	100.0	100.0
Total % known died on waiting list or after transplant	6.1	9.6	11.4	6.4	8.9	10.7
Total % known died or removed as unstable	7.9	11.4	13.2	9.6	12.9	15.1
Total % removed for transplant	60.5	69.3	75.4	60.6	67.6	70.3
Total % with known functioning transplant (alive)	57.9	60.5	36.8	56.7	58.4	39.1

* Follow-up form covering specified time period not yet completed, and possibly has not become due.



B. Waiting List Information

Table B8. Percent of candidates with deceased donor transplants: demographic characteristics
Candidates registered on the waiting list between 07/01/2016 and 06/30/2019

Characteristic	Percent transplanted at time periods since listing									
	This Center					United States				
	N	30 day	1 year	2 years	3 years	N	30 day	1 year	2 years	3 years
All	318	17.9	59.7	65.4	68.6	14,016	20.8	58.2	65.1	67.6
Ethnicity/Race*										
White	173	17.3	57.2	64.7	67.6	8,635	21.1	58.7	65.8	68.3
African-American	68	14.7	58.8	60.3	61.8	3,219	18.9	54.4	61.4	63.7
Hispanic/Latino	50	14.0	58.0	64.0	70.0	1,456	20.5	60.4	67.1	69.6
Asian	25	32.0	80.0	84.0	88.0	538	30.9	67.7	73.2	74.7
Other	2	100.0	100.0	100.0	100.0	168	16.1	57.7	61.3	64.9
Unknown	0	--	--	--	--	0	--	--	--	--
Age										
<2 years	31	25.8	71.0	71.0	71.0	852	13.7	59.9	60.7	60.8
2-11 years	31	29.0	80.6	87.1	90.3	606	12.5	61.6	69.3	71.0
12-17 years	33	63.6	93.9	93.9	93.9	577	32.6	73.5	79.9	81.8
18-34 years	30	13.3	46.7	50.0	53.3	1,347	22.1	56.4	62.1	65.3
35-49 years	52	7.7	48.1	57.7	63.5	2,589	18.3	53.8	61.7	64.7
50-64 years	104	5.8	50.0	59.6	61.5	5,831	20.2	56.6	64.7	67.5
65-69 years	28	14.3	60.7	60.7	67.9	1,849	25.3	61.3	68.0	69.9
70+ years	9	11.1	44.4	44.4	55.6	365	33.4	72.1	73.4	74.0
Gender										
Male	209	14.4	55.5	62.7	66.5	9,873	20.0	56.4	64.0	66.7
Female	109	24.8	67.9	70.6	72.5	4,143	22.9	62.5	67.9	69.7

* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



B. Waiting List Information

Table B9. Percent of candidates with deceased donor transplants: medical characteristics
Candidates registered on the waiting list between 07/01/2016 and 06/30/2019

Characteristic	Percent transplanted at time periods since listing									
	This Center					United States				
	N	30 day	1 year	2 years	3 years	N	30 day	1 year	2 years	3 years
All	318	17.9	59.7	65.4	68.6	14,016	20.8	58.2	65.1	67.6
Blood Type										
O	124	10.5	51.6	53.2	55.6	6,238	13.9	48.8	56.4	59.6
A	126	23.8	59.5	69.0	73.8	5,158	24.9	64.5	71.3	73.2
B	56	17.9	73.2	78.6	80.4	2,003	25.9	64.8	71.1	73.4
AB	12	33.3	83.3	91.7	91.7	617	39.9	78.6	82.5	82.7
Previous Transplant										
Yes	21	23.8	57.1	61.9	61.9	576	17.5	52.1	56.6	58.5
No	297	17.5	59.9	65.7	69.0	13,440	21.0	58.4	65.5	68.0
Primary Disease										
Cardiomyopathy	187	21.9	62.6	69.5	72.7	8,031	22.6	60.4	67.3	69.6
Coronary Artery Disease	50	2.0	40.0	44.0	48.0	3,599	20.3	55.0	63.0	66.2
Retransplant/Graft Failure	18	27.8	61.1	66.7	66.7	497	17.3	53.5	58.1	60.4
Valvular Heart Disease	5	0.0	20.0	20.0	20.0	132	15.9	53.0	59.1	62.1
Congenital Heart Disease	52	15.4	69.2	73.1	75.0	1,536	13.9	57.3	63.2	64.9
Other	6	33.3	83.3	83.3	100.0	221	22.2	49.8	52.9	55.2
Missing	0	--	--	--	--	0	--	--	--	--
Medical Urgency Status at Listing										
Status 1A	119	29.4	75.6	79.0	79.8	3,633	32.5	70.9	73.4	74.2
Status 1B	91	8.8	53.8	63.7	67.0	4,484	15.5	58.9	66.9	69.4
Status 2	49	0.0	40.8	44.9	51.0	2,708	4.8	40.0	51.4	55.5
Unknown	1	100.0	100.0	100.0	100.0	455	8.1	38.7	46.6	50.1



B. Waiting List Information

Table B10. Time to transplant for waiting list candidates*

Candidates registered on the waiting list between 07/01/2016 and 12/31/2021

Percentile	Center	Months to Transplant**		U.S.
		OPO/DSA	Region	
5th	0.2	0.2	0.2	0.2
10th	0.4	0.3	0.3	0.3
25th	1.1	0.8	0.8	0.8
50th (median time to transplant)	4.9	4.4	4.8	4.4
75th	33.8	39.5	Not Observed	Not Observed

* If cells contain "Not Observed" fewer than that percentile of patients had received a transplant. For example, the 50th percentile of time to transplant is the time when 50% of candidates have received transplants. If waiting times are long, then the 50th percentile may not be observed during the follow-up period for this table. Also, if more than 50% of candidates are removed from the list due to death or other reasons before receiving transplants, then the 50th percentile of time to transplant will not be observed.

** Censored on 06/30/2022. Calculated as the months after listing, during which the corresponding percent of all patients initially listed had received a transplant.



B. Waiting List Information

Table B11. Offer Acceptance Practices: 07/01/2021 - 06/30/2022

Offers Acceptance Characteristics	This Center	OPO/DSA	Region	U.S.
Overall				
Number of Offers	3,418	6,624	6,997	60,356
Number of Acceptances	93	253	273	3,338
Expected Acceptances	130.0	274.6	292.6	3,335.3
Offer Acceptance Ratio*	0.72	0.92	0.93	1.00
95% Credible Interval**	[0.58, 0.87]	--	--	--
PHS increased infectious risk				
Number of Offers	910	1,738	1,880	15,775
Number of Acceptances	25	72	77	755
Expected Acceptances	27.3	60.8	67.0	757.5
Offer Acceptance Ratio*	0.92	1.18	1.15	1.00
95% Credible Interval**	[0.61, 1.30]	--	--	--
Ejection fraction < 60				
Number of Offers	1,074	2,162	2,299	19,852
Number of Acceptances	30	75	86	961
Expected Acceptances	36.3	75.6	81.2	950.4
Offer Acceptance Ratio*	0.84	0.99	1.06	1.01
95% Credible Interval**	[0.57, 1.15]	--	--	--
Donor Age >= 40				
Number of Offers	1,615	3,275	3,448	26,961
Number of Acceptances	9	60	62	732
Expected Acceptances	34.4	78.2	83.0	735.0
Offer Acceptance Ratio*	0.30	0.77	0.75	1.00
95% Credible Interval**	[0.15, 0.51]	--	--	--
Hard-to-Place Hearts (Over 50 Offers)				
Number of Offers	1,722	3,208	3,314	22,207
Number of Acceptances	5	26	26	274
Expected Acceptances	20.6	42.9	44.3	278.5
Offer Acceptance Ratio*	0.31	0.62	0.60	0.98
95% Credible Interval**	[0.12, 0.58]	--	--	--
Donor more than 500 miles away				
Number of Offers	1,221	2,081	2,202	20,665
Number of Acceptances	34	57	60	721
Expected Acceptances	31.0	52.9	58.1	758.0
Offer Acceptance Ratio*	1.09	1.08	1.03	0.95
95% Credible Interval**	[0.76, 1.48]	--	--	--

* The offer acceptance ratio estimates the relative offer acceptance practice of NY Presbyterian Hospital/Columbia Univ. Medical Center compared to the national offer acceptance practice. A ratio above one indicates the program is more likely to accept an offer compared to national offer acceptance practices (e.g., an offer acceptance ratio of 1.25 indicates a 25% more likely to accept an offer), while a ratio below one indicates the program is less likely to accept an offer compared to national offer acceptance practices (e.g., an offer acceptance ratio of 0.75 indicates a 25% less likely to accept an offer).

** As an example, the 95% Credible Interval for the overall offer acceptance ratio, [0.58, 0.87], indicates the location of NYCP's true offer acceptance ratio with 95% probability. The best estimate is 28% less likely to accept an offer compared to national acceptance behavior, but NYCP's performance could plausibly range from 42% reduced acceptance up to 13% reduced acceptance.



B. Waiting List Information

Figure B10. Offer acceptance: Overall




Figure B11. Offer acceptance: PHS increased infectious risk




Figure B12. Offer acceptance: Ejection fraction < 60




Figure B13. Offer acceptance: Donor age >= 40




Figure B14. Offer acceptance: Offer number > 50





Figure B15. Offer acceptance: Donor more than 500 miles away





C. Transplant Information

Table C1D. Deceased donor transplant recipient demographic characteristics

Patients transplanted between 07/01/2021 and 06/30/2022

Characteristic	Percentage in each category		
	Center (N=106)	Region (N=325)	U.S. (N=3,813)
Ethnicity/Race (%)*			
White	47.2	48.6	57.0
African-American	19.8	25.5	24.7
Hispanic/Latino	20.8	17.8	12.7
Asian	11.3	7.4	4.6
Other	0.9	0.6	0.9
Unknown	0.0	0.0	0.0
Age (%)			
<2 years	4.7	2.2	3.9
2-11 years	12.3	4.6	3.9
12-17	6.6	3.1	4.5
18-34	5.7	8.0	10.6
35-49 years	15.1	16.3	16.5
50-64 years	36.8	46.2	41.6
65-69 years	14.2	12.9	15.1
70+ years	4.7	6.8	3.9
Gender (%)			
Male	73.6	74.2	72.0
Female	26.4	25.8	28.0

* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



C. Transplant Information

Table C2D. Deceased donor transplant recipient medical characteristics
Patients transplanted between 07/01/2021 and 06/30/2022

Characteristic	Percentage in each category		
	Center (N=106)	Region (N=325)	U.S. (N=3,813)
Blood Type (%)			
O	47.2	46.2	42.3
A	34.0	33.5	38.8
B	15.1	15.7	14.5
AB	3.8	4.6	4.5
Previous Transplant (%)			
Yes	6.6	3.7	3.4
No	93.4	96.3	96.6
Body Mass Index (%)			
0-20	28.3	16.3	16.3
21-25	28.3	31.7	28.6
26-30	19.8	29.5	30.6
31-35	19.8	17.2	19.1
36-40	2.8	3.7	4.1
41+	0.9	1.5	0.8
Unknown	0.0	0.0	0.4
Primary Disease (%)			
Cardiomyopathy	67.9	67.1	62.6
Coronary Artery Disease	14.2	24.0	24.5
Retransplant/Graft Failure	0.0	0.0	0.0
Valvular Heart Disease	0.9	0.9	0.9
Congenital Heart Disease	16.0	6.5	9.9
Other	0.9	1.5	1.9
Missing	0.0	0.0	0.3
Medical Urgency Status at Transplant (%)			
Status 1A	22.6	8.9	10.4
Status 1B	0.9	1.2	2.2
Status 2	0.0	0.0	0.2
Adult Status 1	7.5	9.2	9.7
Adult Status 2	48.1	51.1	44.5
Adult Status 3	6.6	9.5	11.5
Adult Status 4	12.3	15.7	16.2
Adult Status 5	0.0	1.2	0.8
Adult Status 6	1.9	3.1	4.6
Recipient Medical Condition at Transplant (%)			
Not Hospitalized	22.6	26.5	28.2
Hospitalized	15.1	19.4	17.0
ICU	62.3	54.2	54.4
Unknown	0.0	0.0	0.4
Recipient Circulatory Support Status at Transplant (%)			
No Support Mechanism	23.6	18.5	23.1
Devices*	65.1	70.2	61.4
Other Support Mechanism	11.3	11.4	15.4
Unknown	0.0	0.0	0.1

* Devices include ventricular assist devices (VAD), extracorporeal membrane oxygenation (ECMO), intraaortic balloon pump (IABP), and total artificial heart (TAH).



C. Transplant Information

Table C3D. Deceased donor characteristics

Transplants performed between 07/01/2021 and 06/30/2022

Donor Characteristic	Percentage in each category		
	Center (N=106)	Region (N=325)	U.S. (N=3,813)
Cause of Death (%)			
Deceased: Stroke	10.4	13.2	12.2
Deceased: MVA	15.1	14.5	18.7
Deceased: Other	74.5	72.3	69.1
Ethnicity/Race (%)*			
White	69.8	65.8	60.3
African-American	11.3	16.3	17.4
Hispanic/Latino	16.0	15.4	19.2
Asian	2.8	2.2	1.6
Other	0.0	0.3	1.5
Not Reported	0.0	0.0	0.0
Age (%)			
<2 years	2.8	1.5	2.8
2-11 years	10.4	4.3	4.1
12-17	6.6	4.3	6.5
18-34	50.0	50.8	50.7
35-49 years	29.2	33.8	31.5
50-64 years	0.9	5.2	4.4
65-69 years	0.0	0.0	0.0
70+ years	0.0	0.0	0.0
Gender (%)			
Male	50.0	65.2	73.0
Female	50.0	34.8	27.0
Blood Type (%)			
O	51.9	53.8	53.6
A	34.9	31.1	34.5
B	10.4	12.0	10.2
AB	2.8	3.1	1.6
Unknown	0.0	0.0	0.0

* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



C. Transplant Information

Table C4D. Deceased donor transplant characteristics
Transplants performed between 07/01/2021 and 06/30/2022

Transplant Characteristic	Percentage in each category		
	Center (N=106)	Region (N=325)	U.S. (N=3,813)
Total Ischemic Time (Minutes): Local (%)			
Deceased: 0-90 min	6.7	26.1	8.7
Deceased: 91-180 min	20.0	49.3	59.3
Deceased: 181-270 min	40.0	17.4	27.3
Deceased: 271-360 min	26.7	5.8	3.0
Deceased: 361+ min	6.7	1.4	1.3
Not Reported	0.0	0.0	0.4
Total Ischemic Time (Minutes): Shared (%)			
Deceased: 0-90 min	0.0	0.8	0.6
Deceased: 91-180 min	15.4	20.3	13.7
Deceased: 181-270 min	62.6	60.2	67.8
Deceased: 271-360 min	13.2	10.2	13.5
Deceased: 361+ min	5.5	7.4	4.0
Not Reported	3.3	1.2	0.4
Procedure Type (%)			
Single organ	88.7	84.9	88.5
Multi organ	11.3	15.1	11.5
Donor Location (%)			
Local Donation Service Area (DSA)	14.2	21.2	21.5
Another Donation Service Area (DSA)	85.8	78.8	78.5
Median Time in Hospital After Transplant	24.0 Days	18.0 Days	18.0 Days



C. Transplant Information

Table C5D. Adult (18+) 1-month survival with a functioning deceased donor graft

Single organ transplants performed between 07/01/2019 and 03/12/2020, and 06/13/2020 and 12/31/2021

Deaths and retransplants are considered graft failures

Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	136	6,501
Estimated probability of surviving with a functioning graft at 1 month (unadjusted for patient and donor characteristics)	98.51%	96.68%
Expected probability of surviving with a functioning graft at 1 month (adjusted for patient and donor characteristics)	97.19%	--
Number of observed graft failures (including deaths) during the first month after transplant	2	212
Number of expected graft failures (including deaths) during the first month after transplant	3.79	--
Estimated hazard ratio*	0.69	--
95% credible interval for the hazard ratio**	[0.19, 1.51]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.19, 1.51], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 31% lower risk of graft failure compared to an average program, but NYCP's performance could plausibly range from 81% reduced risk up to 51% increased risk.

Figure C1D. Adult (18+) 1-month deceased donor graft failure HR estimate





Figure C2D. Adult (18+) 1-month deceased donor graft failure HR program comparison





C. Transplant Information

Table C6D. Adult (18+) 90-Day survival with a functioning deceased donor graft

Single organ transplants performed between 07/01/2019 and 03/12/2020, and 06/13/2020 and 12/31/2021

Deaths and retransplants are considered graft failures

Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	136	6,501
Estimated probability of surviving with a functioning graft at 90 days (unadjusted for patient and donor characteristics)	96.87%	94.57%
Expected probability of surviving with a functioning graft at 90 days (adjusted for patient and donor characteristics)	95.35%	--
Number of observed graft failures (including deaths) during the first 90 days after transplant	4	340
Number of expected graft failures (including deaths) during the first 90 days after transplant	6.17	--
Estimated hazard ratio*	0.73	--
95% credible interval for the hazard ratio**	[0.27, 1.43]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.27, 1.43], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 27% lower risk of graft failure compared to an average program, but NYCP's performance could plausibly range from 73% reduced risk up to 43% increased risk.

Figure C3D. Adult (18+) 90-Day deceased donor graft failure HR estimate





Figure C4D. Adult (18+) 90-Day deceased donor graft failure HR program comparison





C. Transplant Information

Table C7D. Adult (18+) 1-year survival with a functioning deceased donor graft

Single organ transplants performed between 07/01/2019 and 03/12/2020, and 06/13/2020 and 12/31/2021

Deaths and retransplants are considered graft failures

Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	136	6,501
Estimated probability of surviving with a functioning graft at 1 year (unadjusted for patient and donor characteristics)	92.21%	90.91%
Expected probability of surviving with a functioning graft at 1 year (adjusted for patient and donor characteristics)	92.15%	--
Number of observed graft failures (including deaths) during the first year after transplant	8	501
Number of expected graft failures (including deaths) during the first year after transplant	9.24	--
Estimated hazard ratio*	0.89	--
95% credible interval for the hazard ratio**	[0.43, 1.52]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.43, 1.52], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 11% lower risk of graft failure compared to an average program, but NYCP's performance could plausibly range from 57% reduced risk up to 52% increased risk.

Figure C5D. Adult (18+) 1-year deceased donor graft failure HR estimate





Figure C6D. Adult (18+) 1-year deceased donor graft failure HR program comparison





C. Transplant Information

Table C8D. Adult (18+) 1-year Conditional survival with a functioning deceased donor graft
Single organ transplants performed between 07/01/2019 and 03/12/2020, and 06/13/2020 and 12/31/2021
Deaths and retransplants are considered graft failures
Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	116	5,484
Estimated probability of surviving with a functioning graft at 1 year, among patients with a functioning graft at day 90 (unadjusted for patient and donor characteristics)	95.18%	96.13%
Expected probability of surviving with a functioning graft at 1 year, among patients with a functioning graft at day 90 (adjusted for patient and donor characteristics)	96.65%	--
Number of observed graft failures (including deaths) from day 91 through day 365 after transplant	4	161
Number of expected graft failures (including deaths) from day 91 through day 365 after transplant	3.07	--
Estimated hazard ratio*	1.18	--
95% credible interval for the hazard ratio**	[0.43, 2.30]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.43, 2.30], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 18% higher risk of graft failure compared to an average program, but NYCP's performance could plausibly range from 57% reduced risk up to 130% increased risk.

Figure C7D. Adult (18+) 1-year Conditional deceased donor graft failure HR estimate





Figure C8D. Adult (18+) 1-year Conditional deceased donor graft failure HR program comparison





C. Transplant Information

Table C9D. Adult (18+) 3-year survival with a functioning deceased donor graft

Single organ transplants performed between 01/01/2017 and 06/30/2019

Deaths and retransplants are considered graft failures

Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	138	6,645
Estimated probability of surviving with a functioning graft at 3 years (unadjusted for patient and donor characteristics)	87.29%	85.85%
Expected probability of surviving with a functioning graft at 3 years (adjusted for patient and donor characteristics)	86.55%	--
Number of observed graft failures (including deaths) during the first 3 years after transplant	12	759
Number of expected graft failures (including deaths) during the first 3 years after transplant	14.95	--
Estimated hazard ratio*	0.83	--
95% credible interval for the hazard ratio**	[0.45, 1.31]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.45, 1.31], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 17% lower risk of graft failure compared to an average program, but NYCP's performance could plausibly range from 55% reduced risk up to 31% increased risk.

Figure C9D. Adult (18+) 3-year deceased donor graft failure HR estimate





Figure C10D. Adult (18+) 3-year deceased donor graft failure HR program comparison





C. Transplant Information

Table C10D. Pediatric (<18) 1-month survival with a functioning deceased donor graft
Single organ transplants performed between 07/01/2019 and 03/12/2020, and 06/13/2020 and 12/31/2021
Deaths and retransplants are considered graft failures
Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	51	1,106
Estimated probability of surviving with a functioning graft at 1 month (unadjusted for patient and donor characteristics)	96.08%	97.62%
Expected probability of surviving with a functioning graft at 1 month (adjusted for patient and donor characteristics)	97.48%	--
Number of observed graft failures (including deaths) during the first month after transplant	2	26
Number of expected graft failures (including deaths) during the first month after transplant	1.30	--
Estimated hazard ratio*	1.21	--
95% credible interval for the hazard ratio**	[0.33, 2.66]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.33, 2.66], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 21% higher risk of graft failure compared to an average program, but NYCP's performance could plausibly range from 67% reduced risk up to 166% increased risk.

Figure C11D. Pediatric (<18) 1-month deceased donor graft failure HR estimate





Figure C12D. Pediatric (<18) 1-month deceased donor graft failure HR program comparison





C. Transplant Information

Table C11D. Pediatric (<18) 90-Day survival with a functioning deceased donor graft

Single organ transplants performed between 07/01/2019 and 03/12/2020, and 06/13/2020 and 12/31/2021

Deaths and retransplants are considered graft failures

Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	51	1,106
Estimated probability of surviving with a functioning graft at 90 days (unadjusted for patient and donor characteristics)	92.16%	96.09%
Expected probability of surviving with a functioning graft at 90 days (adjusted for patient and donor characteristics)	95.86%	--
Number of observed graft failures (including deaths) during the first 90 days after transplant	4	42
Number of expected graft failures (including deaths) during the first 90 days after transplant	2.06	--
Estimated hazard ratio*	1.48	--
95% credible interval for the hazard ratio**	[0.54, 2.87]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.54, 2.87], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 48% higher risk of graft failure compared to an average program, but NYCP's performance could plausibly range from 46% reduced risk up to 187% increased risk.

Figure C13D. Pediatric (<18) 90-Day deceased donor graft failure HR estimate





Figure C14D. Pediatric (<18) 90-Day deceased donor graft failure HR program comparison





C. Transplant Information

Table C12D. Pediatric (<18) 1-year survival with a functioning deceased donor graft

Single organ transplants performed between 07/01/2019 and 03/12/2020, and 06/13/2020 and 12/31/2021

Deaths and retransplants are considered graft failures

Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	51	1,106
Estimated probability of surviving with a functioning graft at 1 year (unadjusted for patient and donor characteristics)	87.77%	91.59%
Expected probability of surviving with a functioning graft at 1 year (adjusted for patient and donor characteristics)	91.08%	--
Number of observed graft failures (including deaths) during the first year after transplant	5	74
Number of expected graft failures (including deaths) during the first year after transplant	3.48	--
Estimated hazard ratio*	1.28	--
95% credible interval for the hazard ratio**	[0.51, 2.38]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.51, 2.38], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 28% higher risk of graft failure compared to an average program, but NYCP's performance could plausibly range from 49% reduced risk up to 138% increased risk.

Figure C15D. Pediatric (<18) 1-year deceased donor graft failure HR estimate





Figure C16D. Pediatric (<18) 1-year deceased donor graft failure HR program comparison





C. Transplant Information

Table C13D. Pediatric (<18) 1-year Conditional survival with a functioning deceased donor graft
Single organ transplants performed between 07/01/2019 and 03/12/2020, and 06/13/2020 and 12/31/2021
Deaths and retransplants are considered graft failures
Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	45	957
Estimated probability of surviving with a functioning graft at 1 year, among patients with a functioning graft at day 90 (unadjusted for patient and donor characteristics)	95.24%	95.32%
Expected probability of surviving with a functioning graft at 1 year, among patients with a functioning graft at day 90 (adjusted for patient and donor characteristics)	95.01%	--
Number of observed graft failures (including deaths) from day 91 through day 365 after transplant	1	32
Number of expected graft failures (including deaths) from day 91 through day 365 after transplant	1.42	--
Estimated hazard ratio*	0.88	--
95% credible interval for the hazard ratio**	[0.18, 2.11]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.18, 2.11], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 12% lower risk of graft failure compared to an average program, but NYCP's performance could plausibly range from 82% reduced risk up to 111% increased risk.

Figure C17D. Pediatric (<18) 1-year Conditional deceased donor graft failure HR estimate





Figure C18D. Pediatric (<18) 1-year Conditional deceased donor graft failure HR program comparison





C. Transplant Information

Table C14D. Pediatric (<18) 3-year survival with a functioning deceased donor graft

Single organ transplants performed between 01/01/2017 and 06/30/2019

Deaths and retransplants are considered graft failures

Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	65	1,178
Estimated probability of surviving with a functioning graft at 3 years (unadjusted for patient and donor characteristics)	91.53%	87.18%
Expected probability of surviving with a functioning graft at 3 years (adjusted for patient and donor characteristics)	87.19%	--
Number of observed graft failures (including deaths) during the first 3 years after transplant	5	117
Number of expected graft failures (including deaths) during the first 3 years after transplant	6.69	--
Estimated hazard ratio*	0.81	--
95% credible interval for the hazard ratio**	[0.32, 1.50]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.32, 1.50], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 19% lower risk of graft failure compared to an average program, but NYCP's performance could plausibly range from 68% reduced risk up to 50% increased risk.

Figure C19D. Pediatric (<18) 3-year deceased donor graft failure HR estimate





Figure C20D. Pediatric (<18) 3-year deceased donor graft failure HR program comparison





C. Transplant Information

Table C15D. Adult (18+) 1-month patient survival (deceased donor graft recipients)

Single organ transplants performed between 07/01/2019 and 03/12/2020, and 06/13/2020 and 12/31/2021

Retransplants excluded

Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	131	6,356
Estimated probability of surviving at 1 month (unadjusted for patient and donor characteristics)	100.00%	97.00%
Expected probability of surviving at 1 month (adjusted for patient and donor characteristics)	97.45%	--
Number of observed deaths during the first month after transplant	0	187
Number of expected deaths during the first month after transplant	3.30	--
Estimated hazard ratio*	0.38	--
95% credible interval for the hazard ratio**	[0.05, 1.05]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.05, 1.05], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 62% lower risk of patient death compared to an average program, but NYCP's performance could plausibly range from 95% reduced risk up to 5% increased risk.

Figure C21D. Adult (18+) 1-month patient death HR estimate (deceased donor grafts)





Figure C22D. Adult (18+) 1-month patient death HR program comparison (deceased donor grafts)





C. Transplant Information

Table C16D. Adult (18+) 1-year patient survival (deceased donor graft recipients)

Single organ transplants performed between 07/01/2019 and 03/12/2020, and 06/13/2020 and 12/31/2021

Retransplants excluded

Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	131	6,356
Estimated probability of surviving at 1 year (unadjusted for patient and donor characteristics)	93.47%	91.28%
Expected probability of surviving at 1 year (adjusted for patient and donor characteristics)	92.46%	--
Number of observed deaths during the first year after transplant	6	466
Number of expected deaths during the first year after transplant	8.52	--
Estimated hazard ratio*	0.76	--
95% credible interval for the hazard ratio**	[0.33, 1.37]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.33, 1.37], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 24% lower risk of patient death compared to an average program, but NYCP's performance could plausibly range from 67% reduced risk up to 37% increased risk.

Figure C23D. Adult (18+) 1-year patient death HR estimate (deceased donor grafts)





Figure C24D. Adult (18+) 1-year patient death HR program comparison (deceased donor grafts)





C. Transplant Information

Table C17D. Adult (18+) 3-year patient survival (deceased donor graft recipients)

Single organ transplants performed between 01/01/2017 and 06/30/2019

Retransplants excluded

Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	128	6,474
Estimated probability of surviving at 3 years (unadjusted for patient and donor characteristics)	88.47%	86.58%
Expected probability of surviving at 3 years (adjusted for patient and donor characteristics)	87.37%	--
Number of observed deaths during the first 3 years after transplant	9	705
Number of expected deaths during the first 3 years after transplant	13.20	--
Estimated hazard ratio*	0.72	--
95% credible interval for the hazard ratio**	[0.36, 1.21]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.36, 1.21], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 28% lower risk of patient death compared to an average program, but NYCP's performance could plausibly range from 64% reduced risk up to 21% increased risk.

Figure C25D. Adult (18+) 3-year patient death HR estimate (deceased donor grafts)





Figure C26D. Adult (18+) 3-year patient death HR program comparison (deceased donor grafts)





C. Transplant Information

Table C18D. Pediatric (<18) 1-month patient survival (deceased donor graft recipients)

Single organ transplants performed between 07/01/2019 and 03/12/2020, and 06/13/2020 and 12/31/2021

Retransplants excluded

Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	47	1,057
Estimated probability of surviving at 1 month (unadjusted for patient and donor characteristics)	95.74%	97.60%
Expected probability of surviving at 1 month (adjusted for patient and donor characteristics)	97.22%	--
Number of observed deaths during the first month after transplant	2	25
Number of expected deaths during the first month after transplant	1.32	--
Estimated hazard ratio*	1.20	--
95% credible interval for the hazard ratio**	[0.33, 2.64]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.33, 2.64], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 20% higher risk of patient death compared to an average program, but NYCP's performance could plausibly range from 67% reduced risk up to 164% increased risk.

Figure C27D. Pediatric (<18) 1-month patient death HR estimate (deceased donor grafts)





Figure C28D. Pediatric (<18) 1-month patient death HR program comparison (deceased donor grafts)





C. Transplant Information

Table C19D. Pediatric (<18) 1-year patient survival (deceased donor graft recipients)

Single organ transplants performed between 07/01/2019 and 03/12/2020, and 06/13/2020 and 12/31/2021

Retransplants excluded

Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	47	1,057
Estimated probability of surviving at 1 year (unadjusted for patient and donor characteristics)	86.67%	91.49%
Expected probability of surviving at 1 year (adjusted for patient and donor characteristics)	90.20%	--
Number of observed deaths during the first year after transplant	5	72
Number of expected deaths during the first year after transplant	3.53	--
Estimated hazard ratio*	1.27	--
95% credible interval for the hazard ratio**	[0.51, 2.36]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.51, 2.36], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 27% higher risk of patient death compared to an average program, but NYCP's performance could plausibly range from 49% reduced risk up to 136% increased risk.

Figure C29D. Pediatric (<18) 1-year patient death HR estimate (deceased donor grafts)





Figure C30D. Pediatric (<18) 1-year patient death HR program comparison (deceased donor grafts)





C. Transplant Information

Table C20D. Pediatric (<18) 3-year patient survival (deceased donor graft recipients)

Single organ transplants performed between 01/01/2017 and 06/30/2019

Retransplants excluded

Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	NYCP	U.S.
Number of transplants evaluated	64	1,134
Estimated probability of surviving at 3 years (unadjusted for patient and donor characteristics)	91.38%	87.63%
Expected probability of surviving at 3 years (adjusted for patient and donor characteristics)	87.64%	--
Number of observed deaths during the first 3 years after transplant	5	109
Number of expected deaths during the first 3 years after transplant	6.35	--
Estimated hazard ratio*	0.84	--
95% credible interval for the hazard ratio**	[0.34, 1.56]	--

* The hazard ratio provides an estimate of how NY Presbyterian Hospital/Columbia Univ. Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If NYCP's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

** The 95% credible interval, [0.34, 1.56], indicates the location of NYCP's true hazard ratio with 95% probability. The best estimate is 16% lower risk of patient death compared to an average program, but NYCP's performance could plausibly range from 66% reduced risk up to 56% increased risk.

Figure C31D. Pediatric (<18) 3-year patient death HR estimate (deceased donor grafts)





Figure C32D. Pediatric (<18) 3-year patient death HR program comparison (deceased donor grafts)





C. Transplant Information

Table C21. Multi-organ transplant graft survival: 07/01/2019 - 12/31/2021

Adult (18+) Transplants

Transplant Type	First-Year Outcomes					
	Transplants Performed		Heart Graft Failures		Estimated Heart Graft Survival	
	NYCP-TX1	USA	NYCP-TX1	USA	NYCP-TX1	USA
Heart-Lung	4	116	1	18	75.0%	83.7%
Kidney-Heart	11	736	2	78	81.8%	89.2%
Liver-Heart	4	102	0	16	100.0%	83.8%

Pediatric (<18) Transplants

Transplant Type	First-Year Outcomes					
	Transplants Performed		Heart Graft Failures		Estimated Heart Graft Survival	
	NYCP-TX1	USA	NYCP-TX1	USA	NYCP-TX1	USA
Kidney-Heart	3	13	1	3	66.7%	75.2%

Table C22. Multi-organ transplant patient survival: 07/01/2019 - 12/31/2021

Adult (18+) Transplants

Transplant Type	First-Year Outcomes					
	Transplants Performed		Patient Deaths		Estimated Patient Survival	
	NYCP-TX1	USA	NYCP-TX1	USA	NYCP-TX1	USA
Heart-Lung	4	116	1	18	75.0%	83.7%
Kidney-Heart	11	736	2	77	81.8%	89.4%
Liver-Heart	4	102	0	16	100.0%	83.8%

Pediatric (<18) Transplants

Transplant Type	First-Year Outcomes					
	Transplants Performed		Patient Deaths		Estimated Patient Survival	
	NYCP-TX1	USA	NYCP-TX1	USA	NYCP-TX1	USA
Kidney-Heart	3	13	1	3	66.7%	75.2%