



---

## COVID-19 Guide

---

### Adjustments to Transplant Program and OPO Evaluation Metrics

The Scientific Registry of Transplant Recipients (SRTR), under contract from the Health Resources and Services Administration (HRSA), is charged with evaluating the performance of the nation's transplant system through publication of semi-annual transplant program-specific reports (PSRs) and organ procurement organization (OPO)-specific reports (OSRs). These reports contain performance metrics covering various time periods. For OPOs, these metrics include deceased donor organ yield. For transplant programs, they include pre-transplant mortality rates (formerly called waitlist mortality rates), transplant rates, organ offer acceptance rates, patient mortality after listing, and 1-month, 90-day, 1-year, 1-year conditional on 90-day, and 3-year posttransplant outcomes including graft survival and patient survival.

In response to the current global pandemic, SRTR modified the evaluation metrics for transplant programs and OPOs for the reports released in January 2021, July 2021, January 2022, July 2022, January 2023, July 2023, January 2024, July 2024, January 2025 and July 2025. These reports made adjustments to transplant program and OPO performance metrics so that data during the time around the declaration of a national public health emergency on March 13, 2020, were not included in the metrics.

Modifications for the January 2026 reporting cycle were considered at the Analytic Methods Subcommittee of the SRTR Review Committee (SRC) at its meeting on March 24, 2021, and the full SRC meetings April 27, 2021 and on January 11, 2022. Both the Analytic Methods Subcommittee and the full SRC recommended an ongoing carve out of the first quarter of the pandemic (March 13, 2020 through June 12, 2020) from adjusted performance metrics, as detailed below. These recommendations were reviewed by HRSA's Division of Transplantation, which oversees SRTR. HRSA approved these recommendations, which SRTR will implement for the January 2026 reporting cycle:

Posttransplant Outcomes (including 1-month, 90-day, 1-year, 1-year conditional on 90-day, and 3-year graft and patient survival): Evaluation cohorts will exclude transplants performed between March 13, 2020 and June 12, 2020, inclusive of March 13 and June 12. Patients given transplants before March 13, 2020 will have follow-up censored on March 12, 2020. Patients given transplants after June 12, 2020 will resume normal follow-up. Follow-up will not resume for patients given transplants before March 13, 2020 who are alive with function on June 12, 2020; however, this may be reconsidered as SRTR continues to explore moving to a period-prevalent methodology:

1-month, 90-day, 1-year & 1-year conditional on 90-day Patient and Graft Survival Evaluations: Transplants 7/1/2022-12/31/2024, follow-up through 6/30/2025.

3-year Patient and Graft Survival Evaluations: Transplants 1/1/2020-3/12/2020, follow-up through 3/12/2020. Transplants 6/13/2020-6/30/2022; follow-up through 6/30/2025.

Pre-Transplant Mortality Rate (formerly called Waitlist Mortality Rate): These evaluations are based on normal reporting cohorts.

Days after listing (and before transplant) between 7/1/2023 and 6/30/2025.



---

## COVID-19 Guide

---

Transplant Rate: These evaluations are based on normal reporting cohorts.

Candidates on the waitlist 7/1/2023-6/30/2025.

Overall Rate of Mortality After Listing: These evaluations are based on normal reporting cohorts.

Evaluation period: 7/1/2023-6/30/2025.

Offer Acceptance Rate: These evaluations are based on normal reporting cohorts.

Offers received 7/1/2024-6/30/2025.

These decisions will apply to the evaluations released in the SRTR's semi-annual program-specific reports scheduled for release on January 6, 2026. These changes have been communicated to the leadership of the Organ Procurement and Transplantation Network's (OPTN) Membership and Professional Standards Committee (MSPC).

As with the July 2025 reports, SRTR will continue to report descriptive data beyond March 12, 2020, e.g., waitlist counts, transplant counts, recipient characteristics, donor counts, donor characteristics, etc., but will alter data for performance evaluation metrics as described above.



---

## User Guide

---

This report contains a wide range of useful information about the lung transplant program at CHI St. Luke's Health Baylor College of Medicine Medical Center. The report has three main sections:

- A. Program Summary
- B. Waiting List Information
- C. Transplant Information

The Program Summary is a one-page summary highlighting characteristics of the program, including the number of candidates on the waiting list, the number of transplants performed at the program, the number of patients being cared for by the program, and patient outcomes, including outcomes while on the waiting list (the transplant rate and the death rate while on the waiting list) and outcomes after transplant (patient and graft survival probabilities). If the program performed transplants in both adults and children, survival probabilities for adults and children (pediatrics) are provided separately. For each of the outcomes measures presented, a comparison is provided showing what would be expected at this program if it were performing as similar programs around the country perform when treating similar patients. More details regarding these outcome measures are provided in Sections B and C of the report.

The Waiting List Information section contains more detailed information on how many candidates are on the waiting list at the program, the types of candidates on the waiting list, how long candidates typically have to wait for a transplant at this program, how frequently candidates successfully receive a transplant, and how often candidates on the waiting list die before receiving a transplant.

Table B1 shows the activity on this program's waiting list during two recent 1-year periods and provides comparisons to all programs within this program's OPTN region (see <http://optn.transplant.hrsa.gov/members/regions.asp> for information on OPTN regions) and the nation as a whole. Tables B2 and B3 describe the candidates on the waiting list at this program, with comparisons to candidates waiting in the same donor service area (OPO/DSA) the OPTN region, and the nation as a whole.

Table B4 shows how many candidates were removed from the waiting list because they received a transplant. The program's transplant rate is calculated as the number of candidates who received a transplant divided by the person-years observed at the program (person-years is a combination of how many candidates were on the waiting list along with how long each candidate was followed since some candidates are not on the waiting list for the entire year). The transplant rate and comparisons to what would be expected at this program are presented in Figures B1 and B2. Figure B1 shows the transplant rate compared to what was expected at this program. The expected transplant rate is an estimate of what we would expect at this program if it were performing transplants at rates similar to other programs in the US with similar candidates on their waiting lists. The expected rate is only an estimate, and is made with a certain level of uncertainty. This uncertainty is shown in Figure B2. Figure B2 displays the ratio of the observed to the expected transplant rate. A ratio of 1 indicates that the observed transplant rate was equal to the expected transplant rate, while a ratio less than 1 indicates the observed rate was lower than expected rate and a ratio greater than 1 indicates the observed rate was higher than the expected rate. However, the level of uncertainty must be considered when interpreting these numbers. The 95% interval is also shown on Figure B2. This interval provides a range within which the true ratio of observed to expected transplant rates is likely to be. If this



---

## User Guide

---

confidence interval includes (crosses) 1.0, then we cannot say that this program's observed transplant rate is different from what would be expected. The observed deceased donor transplant rate at this program was 603.6 per 100 person-years. Transplant rates are also provided for adult and pediatric patients separately along with comparisons to adult and pediatric rates in the DSA, the OPTN region, and the nation. Please refer to the PSR Technical Methods documentation available at <http://www.srtr.org> for more detail regarding how expected rates are calculated.

The pre-transplant mortality rate (previously called the waiting list mortality rate) for candidates on the waiting list is presented in Table B5 and Figures B4-B6. These data are presented in the same way as the transplant rate data in the previous section. The intent of this table and figures is to describe risk of death once candidates are listed rather than while they are listed, but before they are transplanted. Therefore, time at risk and deaths after removal from the waiting list for reasons other than transplant, transfer to another transplant program, or recovery (no longer needing a transplant), and before any subsequent transplant, are included. As with transplant rates, mortality rates should be interpreted carefully taking into consideration the interval displayed in Figure B5. For a complete description of how observed and expected mortality rates are calculated, please refer to the technical documentation available at <http://www.srtr.org>.

Survival from listing is presented in Table B6 and Figures B7-B9. These data are presented in the same way as the pre-transplant mortality rate data in the previous section. The intent of this table and figures is to describe risk of death once candidates are listed rather than while they are listed, including after a transplant. As with transplant rates, mortality rates should be interpreted carefully taking into consideration the interval displayed in Figure B8. For a complete description of how observed and expected mortality rates are calculated, please refer to the technical documentation available at <http://www.srtr.org>.

Table B7 presents information on what happens to candidates on the waiting list by three different time points after listing: 6 months, 12 months, and 18 months. The table displays percentages of candidates who have died, been removed from the waiting list, been transplanted, or been transferred or lost-to-follow-up. Tables B8 and B9 provide more detail regarding how many candidates have received a deceased donor transplant by certain time points during the first 3 years after being put on the transplant waiting list. Each row of Tables B8 and B9 presents the percent of candidates who received a deceased donor transplant by each time point. Table B10 presents data on the time it took for different percentages of patients to be transplanted for candidates added to the list between 07/01/2019 and 12/31/2024. The time it took for 5% (the 5th percentile) of patients to receive a transplant at this program was 0.1 months. If "Not Observed" is displayed in the table, then too few candidates received transplants before 06/30/2025 to calculate a particular percentile of transplant times.

Table B11 contains a summary of the offer acceptance practices of the program. The offer acceptance ratio indicates whether the program is more or less likely to accept offers than the average program. If the offer acceptance ratio is greater than 1.0, then the program tends to accept more offers than average; if the offer acceptance ratio is less than 1.0, then the program tends to accept fewer offers than average. Figure B10 shows the distribution of program offer acceptance rates as well as the offer acceptance rate for this program. Figures B11 - B15 similarly show offer acceptance rates for subsets of offers.



---

## User Guide

---

The Transplant Information section begins with descriptions of transplant recipients in Tables C1 and C2. Data on recipients of deceased donor transplants are presented (Tables C1D and C2D); if applicable, data on recipients of living donor transplants are presented separately (Tables C1L and C2L). Comparisons to the region and the nation as a whole are provided. A description of the deceased donors used at this program is provided in Table C3D, along with characteristics of living donors in Table C3L, if applicable. Finally, information on the transplant procedure for deceased and living donor transplants is presented in Tables C4D and C4L, respectively.

Starting with Table C5, transplant outcomes are presented along with comparisons to what would be expected at this program and what happened in the nation as a whole. Tables C5-C14 (tables C5-C10 for Pancreas) present information on graft survival (survival of the transplanted organ), with data presented separately for adult and pediatric recipients. Patients are followed from the time of transplant until either failure of the transplanted organ or death, whichever comes first. Please refer to the technical methods for more information on these calculations (<http://www.srtr.org>).

While Tables C5-C14 present data on graft survival, Tables C15-C20 (tables C11-C20 for Pancreas) present information on patient survival. For these tables, patients are followed from the time of transplant until death, regardless of whether the transplant is functioning or the patient required another transplant to survive.

Tables C21 and C22 summarize the multiorgan transplant outcomes at this program. The summary statistics in these tables are descriptive and are not risk-adjusted for different donor and candidate characteristics.

Additional information regarding the technical methods and the risk adjustment models used to estimate expected event rates is available on the SRTR website at <http://www.srtr.org>. We welcome and encourage feedback on these reports. Please feel free to share feedback with the SRTR at the following e-mail: [srtr@srtr.org](mailto:srtr@srtr.org).




## Table of Contents

Section	Page
COVID-19 Guide	i
User Guide	iii
<b>A. Program Summary</b>	
Program Summary	1
<b>B. Waiting List Information</b>	
Waiting list activity	2
Demographic characteristics of waiting list candidates	3
Medical characteristics of waiting list candidates	4
Deceased donor transplant rates	5
Pre-transplant mortality rates (formerly called Waiting list mortality rates)	6
Patient survival from listing	7
Waiting list candidate status after listing	8
Percent of candidates with deceased donor transplants: demographic characteristics	9
Percent of candidates with deceased donor transplants: medical characteristics	10
Time to transplant for waiting list candidates	11
Offer acceptance practices	12
<b>C. Transplant Information</b>	
Deceased donor transplant recipient demographic characteristics	14
Deceased donor transplant recipient medical characteristics	15
Deceased donor characteristics	16
Deceased donor transplant characteristics	17
Deceased donor graft survival	18
Deceased donor patient survival	28
Multi-organ transplant graft survival	34
Multi-organ transplant patient survival	34



## A. Program Summary

**Figure A1. Waiting list and transplant activity**




**Table A1. Census of transplant recipients**


Recipients	07/01/2023-06/30/2024	07/01/2024-06/30/2025
Transplanted at this center	103	105
Followed by this center*	263	325
...transplanted at this program	249	315
...transplanted elsewhere	14	10

\* Recipients followed are transplant recipients for whom the center has submitted a post-transplant follow-up form for a transplant that took place before the 12-month interval for each column.


**Figure A2. Transplant rates  
07/01/2023 - 06/30/2025**



**Figure A3. Pre-transplant mortality rates  
07/01/2023 - 06/30/2025**



**Figure A4. First-year adult graft and patient survival: 07/01/2022 - 12/31/2024**



**Figure A5. First-year pediatric graft and patient survival: 07/01/2022 - 12/31/2024**

This center did not perform any transplants relevant to this figure during 07/01/2022-12/31/2024



## B. Waiting List Information

**Table B1. Waiting list activity summary: 07/01/2023 - 06/30/2025**

Waiting List Registrations	Counts for this center		Activity for 07/01/2024 to 06/30/2025 as percent of registrants on waiting list on 07/01/2024		
	07/01/2023-06/30/2024	07/01/2024-06/30/2025	This Center (%)	OPTN Region (%)	U.S. (%)
<b>On waiting list at start</b>	16	9	100.0	100.0	100.0
<b>Additions</b>					
New listings at this center	102	122	1,355.6	396.4	411.1
<b>Removals</b>					
Transferred to another center	0	0	0.0	3.6	1.1
Received living donor transplant*	0	0	0.0	0.0	0.0
Received deceased donor transplant*	103	105	1,166.7	351.8	374.3
Died	2	2	22.2	16.1	8.9
Transplanted at another center	0	0	0.0	0.0	2.2
Deteriorated	1	5	55.6	9.8	9.6
Recovered	1	0	0.0	5.4	3.3
Other reasons	2	2	22.2	12.5	14.2
<b>On waiting list at end of period</b>	9	17	188.9	97.3	97.6

\* These patients were removed from waiting list with removal code indicating transplant; this may not equal the number of transplants performed at this center during the specified period.



## B. Waiting List Information

**Table B2. Demographic characteristics of waiting list candidates**

Candidates registered on the waiting list between 07/01/2024 and 06/30/2025

Demographic Characteristic	New Waiting List Registrations 07/01/2024 to 06/30/2025 (%)			All Waiting List Registrations on 06/30/2025 (%)		
	This Center	OPTN Region	U.S.	This Center	OPTN Region	U.S.
	(N=122)	(N=444)	(N=3,774)	(N=17)	(N=109)	(N=896)
<b>All (%)</b>	100.0	100.0	100.0	100.0	100.0	100.0
<b>Ethnicity/Race (%)*</b>						
White	51.6	55.4	66.0	64.7	59.6	64.4
African-American	23.8	12.8	10.8	29.4	12.8	13.6
Hispanic/Latino	19.7	25.7	15.2	0.0	24.8	16.2
Asian	2.5	4.1	4.0	0.0	0.9	3.2
Other	0.0	0.9	1.1	0.0	0.0	0.7
Unknown	2.5	1.1	2.9	5.9	1.8	1.9
<b>Age (%)</b>						
<2 years	0.0	0.5	0.2	0.0	0.9	0.4
2-11 years	0.0	1.1	0.3	0.0	0.0	1.2
12-17 years	0.0	0.7	0.5	0.0	0.9	1.1
18-34 years	3.3	4.1	4.1	5.9	4.6	4.0
35-49 years	14.8	12.4	10.8	5.9	14.7	9.0
50-64 years	45.1	38.3	41.8	41.2	33.9	39.2
65-69 years	23.0	27.0	26.0	17.6	32.1	29.9
70+ years	13.9	16.0	16.4	29.4	12.8	15.1
<b>Gender (%)</b>						
Male	47.5	58.6	61.6	70.6	53.2	54.8
Female	52.5	41.4	38.4	29.4	46.8	45.2

\* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



## B. Waiting List Information

**Table B3. Medical characteristics of waiting list candidates**

Candidates registered on the waiting list between 07/01/2024 and 06/30/2025

Medical Characteristic	New Waiting List Registrations 07/01/2024 to 06/30/2025 (%)			All Waiting List Registrations on 06/30/2025 (%)		
	This Center	OPTN Region	U.S.	This Center	OPTN Region	U.S.
	(N=122)	(N=444)	(N=3,774)	(N=17)	(N=109)	(N=896)
<b>All (%)</b>	100.0	100.0	100.0	100.0	100.0	100.0
<b>Blood Type (%)</b>						
O	49.2	53.2	47.8	52.9	65.1	60.6
A	35.2	34.2	37.4	41.2	27.5	27.5
B	9.8	8.8	11.0	5.9	6.4	10.3
AB	5.7	3.8	3.7	0.0	0.9	1.7
Unknown	0.0	0.0	0.0	0.0	0.0	0.0
<b>Previous Transplant (%)</b>						
Yes	5.7	4.1	3.6	23.5	5.5	4.9
No	94.3	95.9	96.4	76.5	94.5	95.1
Unknown	0.0	0.0	0.0	0.0	0.0	0.0
<b>Primary Disease (%)</b>						
Idiopathic Pulmonary Arterial Hypertension	3.3	2.3	4.0	5.9	6.4	7.1
Cystic Fibrosis	0.0	1.6	1.6	0.0	3.7	1.9
Idiopathic Pulmonary Fibrosis	64.8	67.3	64.7	41.2	56.0	55.0
Emphysema/COPD	15.6	18.2	20.1	23.5	24.8	26.3
Other	16.4	10.6	9.5	29.4	9.2	9.6
Missing	0.0	0.0	0.0	0.0	0.0	0.0



## B. Waiting List Information

Table B4D. Deceased donor transplant rates: 07/01/2023 - 06/30/2025

Waiting List Registrations	This Center	OPO/DSA	Region	U.S.
<b>All Candidates</b>				
Count on waiting list at start*	16	79	143	1,017
Person Years**	34.5	128.0	251.9	1,897.3
Removals for Transplant	208	437	791	6,723
<b>Adult (18+) Candidates</b>				
Count on waiting list at start*	16	78	142	994
Person Years**	34.5	124.6	248.4	1,857.1
Removals for transplant	208	422	776	6,668
<b>Pediatric (&lt;18) Candidates</b>				
Count on waiting list at start*	--	--	--	--
Person Years**	--	--	--	--
Removals for transplant	--	--	--	--

\* Counts in this table may be lower than similar counts in other waiting list tables, such as Table B1. A small percentage (~1%) of patients are found to have died or been transplanted before being removed from the waiting list, so these patients are excluded if the event occurs prior to the start of the study period. Inactive time on the waiting list is included in the calculations for this table.

\*\* Person years are calculated as days (converted to fractional years). The number of days from July 1 or from the date of first wait listing until death, transplant, removal from the waiting list or June 30.

Figure B1D. Observed and expected deceased donor transplant rates: 07/01/2023 - 06/30/2025




Figure B2D. Deceased donor transplant rate ratio estimate





Figure B3D. Observed adult (18+) and pediatric (<18) deceased donor transplant rates: 07/01/2023 - 06/30/2025





## B. Waiting List Information

Table B5. Pre-transplant mortality rates: 07/01/2023 - 06/30/2025

Waiting List Registrations	This Center	OPO/DSA	Region	U.S.
<b>All Candidates</b>				
Count on waiting list at start*	16	79	143	1,017
Person Years**	39.8	146.7	282.3	2,126.2
Number of deaths	6	23	47	301
<b>Adult (18+) Candidates</b>				
Count on waiting list at start*	16	78	142	994
Person Years**	39.8	139.4	275.0	2,078.1
Number of deaths	6	23	47	298
<b>Pediatric (&lt;18) Candidates</b>				
Count on waiting list at start*	--	--	--	--
Person Years**	--	--	--	--
Number of deaths	--	--	--	--

\* Counts in this table may be lower than similar counts in other waiting list tables, such as Table B1. A small percentage (~1%) of patients are found to have died or been transplanted before being removed from the waiting list, so these patients are excluded if the event occurs prior to the start of the study period. Inactive time on the waiting list is included in the calculations for this table.

\*\* Person years are calculated as days (converted to fractional years). The number of days from July 1 or from the date of first wait listing until death, transplant, 60 days after recovery, transfer or June 30.

Figure B4. Observed and expected pre-transplant mortality rates: 07/01/2023 - 06/30/2025




Figure B5. Pre-transplant mortality rate ratio estimate





Figure B6. Observed adult (18+) and pediatric (<18) pre-transplant mortality rates: 07/01/2023 - 06/30/2025





## B. Waiting List Information

Table B6. Rates of patient mortality after listing: 07/01/2023 - 06/30/2025

Waiting List Registrations	This Center	OPO/DSA	Region	U.S.
<b>All Patients</b>				
Count at risk during the evaluation period	451	1,125	2,064	18,536
Person-years*	541.5	1,452.7	2,640.4	23,999.6
Number of Deaths	85	176	300	2,146
<b>Adult (18+) Patients</b>				
Count at risk during the evaluation period	449	1,065	2,004	18,300
Person-years*	538.4	1,373.1	2,560.7	23,689.6
Number of Deaths	84	168	292	2,121
<b>Pediatric (&lt;18) Patients</b>				
Count at risk during the evaluation period	2	60	60	236
Person-years*	3.1	79.7	79.7	310.0
Number of Deaths	1	8	8	25

\* Person-years are calculated as days (converted to fractional years). The number of days from 07/01/2023, or from the date of first wait listing until death, reaching 5 years after listing or June 30, 2025.

\*\* Patient mortality after listing describes the relative survival experience of patients after listing. It depends on many factors, some of which are outside of the control of the transplant program. For example, availability of organs may not be the same in every part of the country.

Figure B7. Observed and expected rates of patient mortality after listing: 07/01/2023 - 06/30/2025




Figure B8. HR estimate of patient mortality after listing





Figure B9. Observed adult (18+) and pediatric (<18) rates of patient mortality after listing: 07/01/2023 - 06/30/2025





## B. Waiting List Information

**Table B7. Waiting list candidate status after listing**  
Candidates registered on waiting list between 01/01/2023 and 12/31/2023

Waiting list status (survival status)	This Center (N=86)			U.S. (N=3,353)		
	Months Since Listing			Months Since Listing		
	6	12	18	6	12	18
<b>Alive on waiting list (%)</b>	9.3	1.2	0.0	14.9	6.3	2.9
<b>Died on the waiting list without transplant (%)</b>	2.3	2.3	2.3	2.3	2.6	2.9
<b>Removed without transplant (%):</b>						
Condition worsened (status unknown)	2.3	2.3	2.3	1.8	2.1	2.3
Condition improved (status unknown)	1.2	1.2	1.2	0.3	0.4	0.6
Refused transplant (status unknown)	1.2	3.5	3.5	0.2	0.4	0.5
Other	1.2	1.2	1.2	2.4	2.9	3.1
<b>Transplant (living or deceased donor) (%):</b>						
Functioning (alive)	75.6	79.1	46.5	73.4	75.2	48.5
Failed-Retransplanted (alive)	0.0	0.0	0.0	0.1	0.2	0.3
Failed-alive not retransplanted	0.0	0.0	0.0	0.0	0.1	0.0
Died	7.0	9.3	20.9	4.1	7.1	10.4
Status Yet Unknown*	0.0	0.0	22.1	0.4	2.5	28.2
<b>Lost or Transferred (status unknown) (%)</b>	0.0	0.0	0.0	0.1	0.1	0.3
<b>TOTAL (%)</b>	100.0	100.0	100.0	100.0	100.0	100.0
Total % known died on waiting list or after transplant	9.3	11.6	23.3	6.4	9.7	13.3
Total % known died or removed as unstable	11.6	14.0	25.6	8.2	11.8	15.6
Total % removed for transplant	82.6	88.4	89.5	78.1	85.1	87.5
Total % with known functioning transplant (alive)	75.6	79.1	46.5	73.4	75.2	48.5

\* Follow-up form covering specified time period not yet completed, and possibly has not become due.



## B. Waiting List Information

**Table B8. Percent of candidates with deceased donor transplants: demographic characteristics**  
Candidates registered on the waiting list between 07/01/2019 and 06/30/2022

Characteristic	Percent transplanted at time periods since listing									
	This Center					United States				
	N	30 day	1 year	2 years	3 years	N	30 day	1 year	2 years	3 years
<b>All</b>	197	54.3	87.8	88.8	88.8	9,035	39.5	79.1	82.9	83.8
<b>Ethnicity/Race*</b>										
White	112	58.0	90.2	91.1	91.1	6,491	40.9	80.7	84.2	85.0
African-American	29	51.7	82.8	86.2	86.2	906	32.2	73.5	79.1	81.0
Hispanic/Latino	49	49.0	85.7	85.7	85.7	1,190	37.8	75.5	80.3	80.9
Asian	7	42.9	85.7	85.7	85.7	362	36.5	73.8	77.1	77.3
Other	0	--	--	--	--	86	43.0	83.7	84.9	86.0
Unknown	0	--	--	--	--	0	--	--	--	--
<b>Age</b>										
<2 years	0	--	--	--	--	33	9.1	57.6	60.6	60.6
2-11 years	0	--	--	--	--	35	17.1	37.1	37.1	40.0
12-17 years	2	0.0	100.0	100.0	100.0	80	38.8	66.2	68.8	68.8
18-34 years	25	44.0	84.0	84.0	84.0	561	33.7	70.6	74.7	76.5
35-49 years	26	57.7	73.1	76.9	76.9	1,043	39.2	74.9	78.6	79.9
50-64 years	86	48.8	86.0	87.2	87.2	4,078	39.0	78.8	83.2	84.2
65-69 years	32	68.8	96.9	96.9	96.9	2,103	39.6	82.3	85.9	86.4
70+ years	26	65.4	100.0	100.0	100.0	1,102	45.7	85.2	87.7	87.8
<b>Gender</b>										
Male	116	64.7	88.8	88.8	88.8	5,359	46.8	84.1	86.5	87.0
Female	81	39.5	86.4	88.9	88.9	3,676	28.9	71.8	77.7	79.2

\* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



## B. Waiting List Information

**Table B9. Percent of candidates with deceased donor transplants: medical characteristics**  
Candidates registered on the waiting list between 07/01/2019 and 06/30/2022

Characteristic	N	Percent transplanted at time periods since listing								
		This Center				United States				
		30 day	1 year	2 years	3 years	N	30 day	1 year	2 years	3 years
<b>All</b>	197	54.3	87.8	88.8	88.8	9,035	39.5	79.1	82.9	83.8
<b>Blood Type</b>										
O	111	54.1	88.3	89.2	89.2	4,293	36.0	76.5	81.0	82.3
A	67	53.7	86.6	88.1	88.1	3,365	44.2	82.8	86.1	86.6
B	13	61.5	92.3	92.3	92.3	1,053	36.6	76.7	79.8	80.2
AB	6	50.0	83.3	83.3	83.3	324	46.0	82.7	85.8	85.8
<b>Previous Transplant</b>										
Yes	14	35.7	78.6	78.6	78.6	365	35.9	72.9	75.9	77.0
No	183	55.7	88.5	89.6	89.6	8,670	39.6	79.3	83.2	84.1
<b>Primary Disease</b>										
Congenital Disease	0	--	--	--	--	0	--	--	--	--
Retransplant/Graft Failure	0	--	--	--	--	0	--	--	--	--
Idiopathic Pulmonary Arterial Hypertension	8	25.0	75.0	75.0	75.0	640	27.0	59.8	64.5	66.2
Cystic Fibrosis	7	57.1	100.0	100.0	100.0	275	34.5	76.0	80.0	81.5
Idiopathic Pulmonary Fibrosis	126	61.1	88.9	90.5	90.5	5,765	42.5	81.8	84.8	85.4
Alpha-1-Antitrypsin Deficiency	0	--	--	--	--	0	--	--	--	--
Emphysema/COPD	34	38.2	88.2	88.2	88.2	1,662	29.1	77.1	84.2	85.7
Other	22	50.0	81.8	81.8	81.8	693	53.0	80.2	82.3	82.5
Missing	0	--	--	--	--	0	--	--	--	--



## B. Waiting List Information

**Table B10. Time to transplant for waiting list candidates\***

Candidates registered on the waiting list between 07/01/2019 and 12/31/2024

Percentile	Center	Months to Transplant**		U.S.
		OPO/DSA	Region	
5th	0.1	0.1	0.2	0.1
10th	0.2	0.2	0.2	0.2
25th	0.4	0.5	0.6	0.4
50th (median time to transplant)	0.9	1.4	1.7	1.4
75th	2.6	6.2	6.9	5.8

\* If cells contain "Not Observed" fewer than that percentile of patients had received a transplant. For example, the 50th percentile of time to transplant is the time when 50% of candidates have received transplants. If waiting times are long, then the 50th percentile may not be observed during the follow-up period for this table. Also, if more than 50% of candidates are removed from the list due to death or other reasons before receiving transplants, then the 50th percentile of time to transplant will not be observed.

\*\* Censored on 06/30/2025. Calculated as the months after listing, during which the corresponding percent of all patients initially listed had received a transplant.



## B. Waiting List Information

Table B11. Offer Acceptance Practices: 07/01/2024 - 06/30/2025

Offers Acceptance Characteristics	This Center	OPO/DSA	Region	U.S.
<b>Overall</b>				
Number of Offers	2,196	3,731	7,454	66,209
Number of Acceptances	103	216	388	3,380
Expected Acceptances	87.3	169.4	397.4	3,380.9
Offer Acceptance Ratio*	1.18	1.27	0.98	1.00
95% Credible Interval**	[0.96, 1.41]	--	--	--
<b>PHS increased infectious risk</b>				
Number of Offers	350	573	1,215	11,672
Number of Acceptances	7	28	51	537
Expected Acceptances	11.2	22.2	57.7	540.8
Offer Acceptance Ratio*	0.68	1.24	0.89	0.99
95% Credible Interval**	[0.31, 1.20]	--	--	--
<b>Donor was current smoker</b>				
Number of Offers	179	292	622	6,493
Number of Acceptances	4	9	20	228
Expected Acceptances	5.6	10.3	24.9	235.8
Offer Acceptance Ratio*	0.78	0.89	0.82	0.97
95% Credible Interval**	[0.29, 1.53]	--	--	--
<b>Donor age &gt;= 55</b>				
Number of Offers	648	827	1,361	14,881
Number of Acceptances	32	33	53	584
Expected Acceptances	18.6	24.6	48.5	569.9
Offer Acceptance Ratio*	1.65	1.31	1.09	1.02
95% Credible Interval**	[1.14, 2.25]	--	--	--
<b>Hard-to-Place Lungs (Over 50 Offers)</b>				
Number of Offers	401	813	1,731	16,488
Number of Acceptances	10	20	36	354
Expected Acceptances	6.5	13.6	39.6	368.4
Offer Acceptance Ratio*	1.41	1.41	0.91	0.96
95% Credible Interval**	[0.73, 2.32]	--	--	--
<b>Donor more than 500 miles away</b>				
Number of Offers	1,505	2,315	4,579	39,234
Number of Acceptances	67	127	218	1,778
Expected Acceptances	54.9	97.9	223.9	1,788.1
Offer Acceptance Ratio*	1.21	1.29	0.97	0.99
95% Credible Interval**	[0.94, 1.51]	--	--	--

\* The offer acceptance ratio estimates the relative offer acceptance practice of CHI St. Luke's Health Baylor College of Medicine Medical Center compared to the national offer acceptance practice. A ratio above one indicates the program accepts more offers compared to national offer acceptance practices (e.g., an offer acceptance ratio of 1.25 indicates a center accepts 25% more offers than is expected based on national offer acceptance practices), while a ratio below one indicates the program accepts fewer offers compared to national offer acceptance practices (e.g., an offer acceptance ratio of 0.75 indicates a center accepts 25% fewer offers than is expected based on national offer acceptance practices).

\*\* As an example, the 95% Credible Interval for the overall offer acceptance ratio, [0.96, 1.41], indicates the location of TXHI's true offer acceptance ratio with 95% probability. The best estimate is 18% more likely to accept an offer compared to national acceptance behavior, but TXHI's performance could plausibly range from 4% reduced acceptance up to 41% higher acceptance.



## B. Waiting List Information

Figure B10. Offer acceptance: Overall




Figure B11. Offer acceptance: PHS increased infectious risk




Figure B12. Offer acceptance: Donor was current smoker




Figure B13. Offer acceptance: Donor age >= 55




Figure B14. Offer acceptance: Offer number > 50





Figure B15. Offer acceptance: Donor more than 500 miles away





## C. Transplant Information

**Table C1D. Deceased donor transplant recipient demographic characteristics**  
Patients transplanted between 07/01/2024 and 06/30/2025

Characteristic	Percentage in each category		
	Center (N=105)	Region (N=394)	U.S. (N=3,436)
<b>Ethnicity/Race (%)*</b>			
White	49.5	53.8	67.5
African-American	21.9	13.7	10.4
Hispanic/Latino	23.8	25.6	14.2
Asian	2.9	4.8	3.9
Other	0.0	1.3	1.1
Unknown	1.9	0.8	2.9
<b>Age (%)</b>			
<2 years	0.0	0.3	0.0
2-11 years	0.0	1.0	0.3
12-17	0.0	0.5	0.4
18-34	4.8	4.6	4.2
35-49 years	16.2	12.2	10.8
50-64 years	41.9	38.1	41.5
65-69 years	21.9	26.1	25.7
70+ years	15.2	17.3	17.1
<b>Gender (%)</b>			
Male	44.8	56.9	60.7
Female	55.2	43.1	39.3

\* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



## C. Transplant Information

**Table C2D. Deceased donor transplant recipient medical characteristics**  
Patients transplanted between 07/01/2024 and 06/30/2025

Characteristic	Percentage in each category		
	Center (N=105)	Region (N=394)	U.S. (N=3,436)
<b>Blood Type (%)</b>			
O	46.7	51.3	46.7
A	37.1	35.0	38.2
B	9.5	9.6	11.3
AB	6.7	4.1	3.8
<b>Previous Transplant (%)</b>			
Yes	4.8	4.3	3.7
No	95.2	95.7	96.3
<b>Body Mass Index (%)</b>			
0-20	14.3	16.2	14.1
21-25	26.7	34.3	32.8
26-30	31.4	34.8	37.2
31-35	26.7	14.2	14.9
36-40	1.0	0.5	0.7
41+	0.0	0.0	0.2
Unknown	0.0	0.0	0.2
<b>Primary Disease (%)</b>			
Idiopathic Pulmonary Arterial Hypertension	4.8	3.3	3.6
Cystic Fibrosis	1.0	2.0	2.4
Idiopathic Pulmonary Fibrosis	61.9	66.2	63.7
Emphysema/COPD	16.2	17.8	20.5
Other	16.2	10.7	9.7
Missing	0.0	0.0	0.1
<b>Recipient Medical Condition at Transplant (%)</b>			
Not Hospitalized	68.6	60.2	62.6
Hospitalized	1.9	16.8	16.9
ICU	29.5	23.1	20.5
Unknown	0.0	0.0	0.0



## C. Transplant Information

**Table C3D. Deceased donor characteristics**  
Transplants performed between 07/01/2024 and 06/30/2025

Donor Characteristic	Percentage in each category		
	Center (N=105)	Region (N=394)	U.S. (N=3,436)
<b>Cause of Death (%)</b>			
Deceased: Stroke	38.1	29.9	29.5
Deceased: MVA	17.1	16.8	12.3
Deceased: Other	44.8	53.3	58.2
<b>Ethnicity/Race (%)*</b>			
White	47.6	51.0	57.0
African-American	19.0	18.5	17.4
Hispanic/Latino	24.8	24.1	19.5
Asian	5.7	3.3	4.0
Other	1.0	1.5	1.3
Not Reported	1.9	1.5	0.8
<b>Age (%)</b>			
<2 years	0.0	0.3	0.1
2-11 years	0.0	1.0	0.4
12-17	5.7	5.8	4.9
18-34	26.7	36.0	33.9
35-49 years	25.7	35.0	35.6
50-64 years	30.5	18.5	22.0
65-69 years	10.5	3.0	2.4
70+ years	1.0	0.3	0.7
<b>Gender (%)</b>			
Male	64.8	59.4	60.2
Female	35.2	40.6	39.8
<b>Blood Type (%)</b>			
O	57.1	58.9	55.6
A	32.4	31.5	34.1
B	5.7	7.6	9.1
AB	4.8	2.0	1.2
Unknown	0.0	0.0	0.0

\* Race and ethnicity are reported together as a single data element, reflecting their data collection (either race or ethnicity is required, but not both). Patients formerly coded as white and Hispanic are coded as Hispanic. Race and ethnicity sum to 100%.



## C. Transplant Information

**Table C4D. Deceased donor transplant characteristics**  
Transplants performed between 07/01/2024 and 06/30/2025

Transplant Characteristic	Percentage in each category		
	Center (N=105)	Region (N=394)	U.S. (N=3,436)
<b>Total Ischemic Time (Minutes): Local (%)</b>			
Deceased: 0-90 min	0.0	0.0	0.4
Deceased: 91-180 min	0.0	3.8	2.6
Deceased: 181-270 min	20.0	13.2	17.1
Deceased: 271-360 min	46.7	30.2	24.2
Deceased: 361+ min	33.3	52.8	55.8
Not Reported	0.0	0.0	0.0
<b>Total Ischemic Time (Minutes): Shared (%)</b>			
Deceased: 0-90 min	0.0	1.8	0.9
Deceased: 91-180 min	0.0	0.0	0.4
Deceased: 181-270 min	0.0	2.1	2.8
Deceased: 271-360 min	11.1	11.7	14.7
Deceased: 361+ min	88.9	84.5	81.2
Not Reported	0.0	0.0	0.0
<b>Procedure Type (%)</b>			
Single organ	99.0	98.7	98.7
Multi organ	1.0	1.3	1.3
<b>Donor Location (%)</b>			
Local Donation Service Area (DSA)	14.3	13.5	14.7
Another Donation Service Area (DSA)	85.7	86.5	85.3
<b>Median Time in Hospital After Transplant</b>	33.0 Days	21.5 Days	20.0 Days



## C. Transplant Information

**Table C5D. Adult (18+) 1-month survival with a functioning deceased donor graft**

Single organ transplants performed between 07/01/2022 and 12/31/2024


Deaths and retransplants are considered graft failures

	TXHI	U.S.
Number of transplants evaluated	207	7,577
Estimated probability of surviving with a functioning graft at 1 month & [95% CI] (unadjusted for patient and donor characteristics)	98.07% [96.21%-99.96%]	97.58% [97.24%-97.93%]
Expected probability of surviving with a functioning graft at 1 month (adjusted for patient and donor characteristics)	97.25%	--
Number of observed graft failures (including deaths) during the first month after transplant	4	183
Number of expected graft failures (including deaths) during the first month after transplant	5.76	--
Estimated hazard ratio*	0.77	--
95% credible interval for the hazard ratio**	[0.28, 1.50]	--


\* The hazard ratio provides an estimate of how CHI St. Luke's Health Baylor College of Medicine Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If TXHI's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

\*\* The 95% credible interval, [0.28, 1.50], indicates the location of TXHI's true hazard ratio with 95% probability. The best estimate is 23% lower risk of graft failure compared to an average program, but TXHI's performance could plausibly range from 72% reduced risk up to 50% increased risk.

**Figure C1D. Adult (18+) 1-month deceased donor graft failure HR estimate**



**Figure C2D. Adult (18+) 1-month deceased donor graft failure HR program comparison**





## C. Transplant Information

**Table C6D. Adult (18+) 90-Day survival with a functioning deceased donor graft**

Single organ transplants performed between 07/01/2022 and 12/31/2024


Deaths and retransplants are considered graft failures

	TXHI	U.S.
Number of transplants evaluated	207	7,577
Estimated probability of surviving with a functioning graft at 90 days & [95% CI] (unadjusted for patient and donor characteristics)	96.14% [93.54%-98.80%]	95.50% [95.03%-95.97%]
Expected probability of surviving with a functioning graft at 90 days (adjusted for patient and donor characteristics)	94.87%	--
Number of observed graft failures (including deaths) during the first 90 days after transplant	8	341
Number of expected graft failures (including deaths) during the first 90 days after transplant	10.75	--
Estimated hazard ratio*	0.78	--
95% credible interval for the hazard ratio**	[0.38, 1.34]	--


\* The hazard ratio provides an estimate of how CHI St. Luke's Health Baylor College of Medicine Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If TXHI's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

\*\* The 95% credible interval, [0.38, 1.34], indicates the location of TXHI's true hazard ratio with 95% probability. The best estimate is 22% lower risk of graft failure compared to an average program, but TXHI's performance could plausibly range from 62% reduced risk up to 34% increased risk.

**Figure C3D. Adult (18+) 90-Day deceased donor graft failure HR estimate**



**Figure C4D. Adult (18+) 90-Day deceased donor graft failure HR program comparison**





## C. Transplant Information


**Table C7D. Adult (18+) 1-year survival with a functioning deceased donor graft**  
Single organ transplants performed between 07/01/2022 and 12/31/2024  
Deaths and retransplants are considered graft failures

	TXHI	U.S.
Number of transplants evaluated	207	7,577
Estimated probability of surviving with a functioning graft at 1 year & [95% CI] (unadjusted for patient and donor characteristics)	86.47% [81.71%-91.50%]	89.63% [88.91%-90.35%]
Expected probability of surviving with a functioning graft at 1 year (adjusted for patient and donor characteristics)	88.18%	--
Number of observed graft failures (including deaths) during the first year after transplant	26	725
Number of expected graft failures (including deaths) during the first year after transplant	22.44	--
Estimated hazard ratio*	1.15	--
95% credible interval for the hazard ratio**	[0.76, 1.61]	--


\* The hazard ratio provides an estimate of how CHI St. Luke's Health Baylor College of Medicine Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If TXHI's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

\*\* The 95% credible interval, [0.76, 1.61], indicates the location of TXHI's true hazard ratio with 95% probability. The best estimate is 15% higher risk of graft failure compared to an average program, but TXHI's performance could plausibly range from 24% reduced risk up to 61% increased risk.

**Figure C5D. Adult (18+) 1-year deceased donor graft failure HR estimate**



**Figure C6D. Adult (18+) 1-year deceased donor graft failure HR program comparison**





## C. Transplant Information


**Table C8D. Adult (18+) 1-year Conditional survival with a functioning deceased donor graft**  
Single organ transplants performed between 07/01/2022 and 12/31/2024  
Deaths and retransplants are considered graft failures

	TXHI	U.S.
Number of transplants evaluated	199	7,235
Estimated probability of surviving with a functioning graft at 1 year, among patients with a functioning graft at day 90 & [95% CI] (unadjusted for patient and donor characteristics)	89.94% [87.35%-92.61%]	93.85% [93.56%-94.15%]
Expected probability of surviving with a functioning graft at 1 year, among patients with a functioning graft at day 90 (adjusted for patient and donor characteristics)	92.95%	--
Number of observed graft failures (including deaths) from day 91 through day 365 after transplant	18	384
Number of expected graft failures (including deaths) from day 91 through day 365 after transplant	11.69	--
Estimated hazard ratio*	1.46	--
95% credible interval for the hazard ratio**	[0.89, 2.17]	--


\* The hazard ratio provides an estimate of how CHI St. Luke's Health Baylor College of Medicine Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If TXHI's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

\*\* The 95% credible interval, [0.89, 2.17], indicates the location of TXHI's true hazard ratio with 95% probability. The best estimate is 46% higher risk of graft failure compared to an average program, but TXHI's performance could plausibly range from 11% reduced risk up to 117% increased risk.

**Figure C7D. Adult (18+) 1-year Conditional deceased donor graft failure HR estimate**



**Figure C8D. Adult (18+) 1-year Conditional deceased donor graft failure HR program comparison**





## C. Transplant Information

**Table C9D. Adult (18+) 3-year survival with a functioning deceased donor graft**

Single organ transplants performed between 01/01/2020 and 03/12/2020, and 06/13/2020 and 06/30/2022

Deaths and retransplants are considered graft failures


Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	TXHI	U.S.
Number of transplants evaluated	134	5,709
Estimated probability of surviving with a functioning graft at 3 years & [95% CI] (unadjusted for patient and donor characteristics)	67.33% [59.59%-76.08%]	72.69% [71.48%-73.92%]
Expected probability of surviving with a functioning graft at 3 years (adjusted for patient and donor characteristics)	71.78%	--
Number of observed graft failures (including deaths) during the first 3 years after transplant	41	1,409
Number of expected graft failures (including deaths) during the first 3 years after transplant	34.38	--
Estimated hazard ratio*	1.18	--
95% credible interval for the hazard ratio**	[0.86, 1.56]	--


\* The hazard ratio provides an estimate of how CHI St. Luke's Health Baylor College of Medicine Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If TXHI's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

\*\* The 95% credible interval, [0.86, 1.56], indicates the location of TXHI's true hazard ratio with 95% probability. The best estimate is 18% higher risk of graft failure compared to an average program, but TXHI's performance could plausibly range from 14% reduced risk up to 56% increased risk.

**Figure C9D. Adult (18+) 3-year deceased donor graft failure HR estimate**



**Figure C10D. Adult (18+) 3-year deceased donor graft failure HR program comparison**





## C. Transplant Information

**Table C10D. Pediatric (<18) 1-month survival with a functioning deceased donor graft**  
Single organ transplants performed between 07/01/2022 and 12/31/2024  
Deaths and retransplants are considered graft failures

This center did not perform any  
transplants relevant to  
this table during  
07/01/2022-12/31/2024

**Figure C11D. Pediatric (<18)  
1-month deceased donor graft  
failure HR estimate**

This center did not perform any  
transplants relevant to  
this figure during  
07/01/2022-12/31/2024

**Figure C12D. Pediatric (<18) 1-month deceased donor graft  
failure HR program comparison**

This center did not perform any  
transplants relevant to  
this figure during  
07/01/2022-12/31/2024



## C. Transplant Information

### Table C11D. Pediatric (<18) 90-Day survival with a functioning deceased donor graft

Single organ transplants performed between 07/01/2022 and 12/31/2024

Deaths and retransplants are considered graft failures

This center did not perform any  
transplants relevant to  
this table during  
07/01/2022-12/31/2024

### Figure C13D. Pediatric (<18) 90-Day deceased donor graft failure HR estimate

This center did not perform any  
transplants relevant to  
this figure during  
07/01/2022-12/31/2024

### Figure C14D. Pediatric (<18) 90-Day deceased donor graft failure HR program comparison

This center did not perform any  
transplants relevant to  
this figure during  
07/01/2022-12/31/2024



## C. Transplant Information

**Table C12D. Pediatric (<18) 1-year survival with a functioning deceased donor graft**  
Single organ transplants performed between 07/01/2022 and 12/31/2024  
Deaths and retransplants are considered graft failures

This center did not perform any  
transplants relevant to  
this table during  
07/01/2022-12/31/2024

**Figure C15D. Pediatric (<18)  
1-year deceased donor graft  
failure HR estimate**

This center did not perform any  
transplants relevant to  
this figure during  
07/01/2022-12/31/2024

**Figure C16D. Pediatric (<18) 1-year deceased donor graft  
failure HR program comparison**

This center did not perform any  
transplants relevant to  
this figure during  
07/01/2022-12/31/2024



## C. Transplant Information

**Table C13D. Pediatric (<18) 1-year Conditional survival with a functioning deceased donor graft**  
Single organ transplants performed between 07/01/2022 and 12/31/2024  
Deaths and retransplants are considered graft failures

This center did not perform any  
transplants relevant to  
this table during  
07/01/2022-12/31/2024

**Figure C17D. Pediatric (<18)  
1-year Conditional deceased  
donor graft failure HR estimate**

This center did not perform any  
transplants relevant to  
this figure during  
07/01/2022-12/31/2024

**Figure C18D. Pediatric (<18) 1-year Conditional deceased  
donor graft failure HR program comparison**

This center did not perform any  
transplants relevant to  
this figure during  
07/01/2022-12/31/2024



## C. Transplant Information


**Table C14D. Pediatric (<18) 3-year survival with a functioning deceased donor graft**  
Single organ transplants performed between 01/01/2020 and 03/12/2020, and 06/13/2020 and 06/30/2022  
Deaths and retransplants are considered graft failures  
Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	TXHI	U.S.
Number of transplants evaluated	2	61
Estimated probability of surviving with a functioning graft at 3 years & [95% CI] (unadjusted for patient and donor characteristics)	50.00% [12.50%-100.00%]	70.13% [58.74%-83.72%]
Expected probability of surviving with a functioning graft at 3 years (adjusted for patient and donor characteristics)	69.61%	--
Number of observed graft failures (including deaths) during the first 3 years after transplant	1	16
Number of expected graft failures (including deaths) during the first 3 years after transplant	0.48	--
Estimated hazard ratio*	1.21	--
95% credible interval for the hazard ratio**	[0.25, 2.91]	--


\* The hazard ratio provides an estimate of how CHI St. Luke's Health Baylor College of Medicine Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected graft failure rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected graft failure rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If TXHI's graft failure rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

\*\* The 95% credible interval, [0.25, 2.91], indicates the location of TXHI's true hazard ratio with 95% probability. The best estimate is 21% higher risk of graft failure compared to an average program, but TXHI's performance could plausibly range from 75% reduced risk up to 191% increased risk.

**Figure C19D. Pediatric (<18) 3-year deceased donor graft failure HR estimate**



**Figure C20D. Pediatric (<18) 3-year deceased donor graft failure HR program comparison**





## C. Transplant Information

**Table C15D. Adult (18+) 1-month patient survival (deceased donor graft recipients)**


Single organ transplants performed between 07/01/2022 and 12/31/2024  
Retransplants excluded

	TXHI	U.S.
Number of transplants evaluated	187	7,339
Estimated probability of surviving at 1 month & [95% CI] (unadjusted for patient and donor characteristics)	98.40% [96.61%-100.00%]	98.00% [97.68%-98.32%]
Expected probability of surviving at 1 month (adjusted for patient and donor characteristics)	97.77%	--
Number of observed deaths during the first month after transplant	3	147
Number of expected deaths during the first month after transplant	4.20	--
Estimated hazard ratio*	0.81	--
95% credible interval for the hazard ratio**	[0.26, 1.65]	--


\* The hazard ratio provides an estimate of how CHI St. Luke's Health Baylor College of Medicine Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If TXHI's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

\*\* The 95% credible interval, [0.26, 1.65], indicates the location of TXHI's true hazard ratio with 95% probability. The best estimate is 19% lower risk of patient death compared to an average program, but TXHI's performance could plausibly range from 74% reduced risk up to 65% increased risk.

**Figure C21D. Adult (18+) 1-month patient death HR estimate (deceased donor grafts)**



**Figure C22D. Adult (18+) 1-month patient death HR program comparison (deceased donor grafts)**





## C. Transplant Information

**Table C16D. Adult (18+) 1-year patient survival (deceased donor graft recipients)**


Single organ transplants performed between 07/01/2022 and 12/31/2024  
Retransplants excluded

	TXHI	U.S.
Number of transplants evaluated	187	7,339
Estimated probability of surviving at 1 year & [95% CI] (unadjusted for patient and donor characteristics)	88.47% [83.79%-93.41%]	90.25% [89.54%-90.96%]
Expected probability of surviving at 1 year (adjusted for patient and donor characteristics)	89.11%	--
Number of observed deaths during the first year after transplant	20	662
Number of expected deaths during the first year after transplant	18.35	--
Estimated hazard ratio*	1.08	--
95% credible interval for the hazard ratio**	[0.68, 1.58]	--


\* The hazard ratio provides an estimate of how CHI St. Luke's Health Baylor College of Medicine Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If TXHI's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

\*\* The 95% credible interval, [0.68, 1.58], indicates the location of TXHI's true hazard ratio with 95% probability. The best estimate is 8% higher risk of patient death compared to an average program, but TXHI's performance could plausibly range from 32% reduced risk up to 58% increased risk.

**Figure C23D. Adult (18+) 1-year patient death HR estimate (deceased donor grafts)**



**Figure C24D. Adult (18+) 1-year patient death HR program comparison (deceased donor grafts)**





## C. Transplant Information

**Table C17D. Adult (18+) 3-year patient survival (deceased donor graft recipients)**

Single organ transplants performed between 01/01/2020 and 03/12/2020, and 06/13/2020 and 06/30/2022

Retransplants excluded


Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	TXHI	U.S.
Number of transplants evaluated	125	5,530
Estimated probability of surviving at 3 years & [95% CI] (unadjusted for patient and donor characteristics)	70.99% [63.23%-79.71%]	74.12% [72.92%-75.35%]
Expected probability of surviving at 3 years (adjusted for patient and donor characteristics)	73.73%	--
Number of observed deaths during the first 3 years after transplant	34	1,301
Number of expected deaths during the first 3 years after transplant	30.78	--
Estimated hazard ratio*	1.10	--
95% credible interval for the hazard ratio**	[0.77, 1.49]	--


\* The hazard ratio provides an estimate of how CHI St. Luke's Health Baylor College of Medicine Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If TXHI's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

\*\* The 95% credible interval, [0.77, 1.49], indicates the location of TXHI's true hazard ratio with 95% probability. The best estimate is 10% higher risk of patient death compared to an average program, but TXHI's performance could plausibly range from 23% reduced risk up to 49% increased risk.

**Figure C25D. Adult (18+) 3-year patient death HR estimate (deceased donor grafts)**



**Figure C26D. Adult (18+) 3-year patient death HR program comparison (deceased donor grafts)**





## C. Transplant Information

**Table C18D. Pediatric (<18) 1-month patient survival (deceased donor graft recipients)**  
Single organ transplants performed between 07/01/2022 and 12/31/2024  
Retransplants excluded

This center did not perform any  
transplants relevant to  
this table during  
07/01/2022-12/31/2024

**Figure C27D. Pediatric (<18)  
1-month patient death HR  
estimate (deceased donor grafts)**

This center did not perform any  
transplants relevant to  
this figure during  
07/01/2022-12/31/2024

**Figure C28D. Pediatric (<18) 1-month patient death HR  
program comparison (deceased donor grafts)**

This center did not perform any  
transplants relevant to  
this figure during  
07/01/2022-12/31/2024



## C. Transplant Information

**Table C19D. Pediatric (<18) 1-year patient survival (deceased donor graft recipients)**  
Single organ transplants performed between 07/01/2022 and 12/31/2024  
Retransplants excluded

This center did not perform any  
transplants relevant to  
this table during  
07/01/2022-12/31/2024

**Figure C29D. Pediatric (<18)  
1-year patient death HR estimate  
(deceased donor grafts)**

This center did not perform any  
transplants relevant to  
this figure during  
07/01/2022-12/31/2024

**Figure C30D. Pediatric (<18) 1-year patient death HR program  
comparison (deceased donor grafts)**

This center did not perform any  
transplants relevant to  
this figure during  
07/01/2022-12/31/2024



## C. Transplant Information

**Table C20D. Pediatric (<18) 3-year patient survival (deceased donor graft recipients)**

Single organ transplants performed between 01/01/2020 and 03/12/2020, and 06/13/2020 and 06/30/2022  
Retransplants excluded


Follow-up ends on 3/12/2020 for recipients transplanted prior to 3/13/2020

	TXHI	U.S.
Number of transplants evaluated	2	60
Estimated probability of surviving at 3 years & [95% CI] (unadjusted for patient and donor characteristics)	100.00% [100.00%-100.00%]	74.68% [64.02%-87.12%]
Expected probability of surviving at 3 years (adjusted for patient and donor characteristics)	74.91%	--
Number of observed deaths during the first 3 years after transplant	0	14
Number of expected deaths during the first 3 years after transplant	0.58	--
Estimated hazard ratio*	0.78	--
95% credible interval for the hazard ratio**	[0.09, 2.16]	--


\* The hazard ratio provides an estimate of how CHI St. Luke's Health Baylor College of Medicine Medical Center's results compare with what was expected based on modeling the transplant outcomes from all U.S. programs. A ratio above 1 indicates higher than expected patient death rates (e.g., a hazard ratio of 1.5 would indicate 50% higher risk), and a ratio below 1 indicates lower than expected patient death rates (e.g., a hazard ratio of 0.75 would indicate 25% lower risk). If TXHI's patient death rate were precisely the expected rate, the estimated hazard ratio would be 1.0.

\*\* The 95% credible interval, [0.09, 2.16], indicates the location of TXHI's true hazard ratio with 95% probability. The best estimate is 22% lower risk of patient death compared to an average program, but TXHI's performance could plausibly range from 91% reduced risk up to 116% increased risk.

**Figure C31D. Pediatric (<18) 3-year patient death HR estimate (deceased donor grafts)**



**Figure C32D. Pediatric (<18) 3-year patient death HR program comparison (deceased donor grafts)**





## C. Transplant Information

Table C21. Multi-organ transplant graft survival: 07/01/2022 - 12/31/2024

### Adult (18+) Transplants

Transplant Type	First-Year Outcomes					
	Transplants Performed		Lung Graft Failures		Estimated Lung Graft Survival	
	TXHI-TX1	USA	TXHI-TX1	USA	TXHI-TX1	USA
Heart-Lung	1	132	0	19	100.0%	85.6%
Liver-Lung	1	54	1	7	0.0%	86.1%

### Pediatric (<18) Transplants

No pediatric (<18) multi-organ transplants were performed

Table C22. Multi-organ transplant patient survival: 07/01/2022 - 12/31/2024

### Adult (18+) Transplants

Transplant Type	First-Year Outcomes					
	Transplants Performed		Patient Deaths		Estimated Patient Survival	
	TXHI-TX1	USA	TXHI-TX1	USA	TXHI-TX1	USA
Heart-Lung	1	132	0	19	100.0%	85.6%
Liver-Lung	1	54	1	7	0.0%	87.0%

### Pediatric (<18) Transplants

No pediatric (<18) multi-organ transplants were performed